

UNIT – 1

MATRICES AND DETERMINANTS

Unit Outlines

- 1.1 Introduction to Matrices
- 1.2 Types of Matrices
- 1.3 Addition and Subtraction of Matrices
- 1.4 Multiplication of Matrices
- 1.5 Multiplicative inverse of a Matrix
- 1.6 Solution of Simultaneous Linear Equations.

After studying this unit the students will be able to:

- A matrix with real entries and relate its rectangular layout (Formation) with real life.
- Rows and columns of a matrix.
The order of a matrix.
- Equality of two matrices.
- Define and identify row matrix, column matrix, rectangular matrix, square matrix, zero/null matrix, identity matrix, scalar matrix, diagonal matrix, transpose of a matrix, symmetric and skew-symmetric matrices.
- Know whether the given matrices are conformable for addition/subtraction.
- Add and subtract matrices.
- Multiply a matrix by a real number.
- Verify commutative and associative laws under addition.
- Define additive identity of a matrix.
- Known whether the given matrices are conformable for multiplication.
- Multiply two (or three) matrices.
- Verify associative law under multiplication.

- Verify distributive laws.
- Show with the help of an example that commutative law under multiplication does not hold in general (i.e., $AB \neq BA$).
- Define multiplicative identity of a matrix.
- Verify the result $(AB)^t = B^t A^t$.
- Define the Determinant of a square matrix.
- Evaluate determinant of a matrix.
- Define singular and non-singular matrices.
- Define adjoint of a matrix.
- Find multiplicative inverse of a non-singular matrix A and verify that $AA^{-1} = I = A^{-1}A$ where I is the identity matrix.
- Use adjoint method to calculate inverse of a non-singular matrix.
- Verify the result $(AB)^{-1} = B^{-1}A^{-1}$
- Solve a system of two linear equations and related real life problems in two unknowns using
- Matrix inversion method,
- Cramer's rule.

Introduction of Matrix:

The idea of Matrix was given by “Arthur Cayley”, an English mathematician of 19th century. Who first developed “Theory of Matrices” in 1858.

Matrix:

A rectangular array or a formation of collection of real numbers, say 0, 1, 2, 3, 4 and 9, such as; $\begin{bmatrix} 1 & 3 & 4 \\ 9 & 2 & 0 \end{bmatrix}$ and then enclosed by brackets [] is said to form a matrix $\begin{bmatrix} 1 & 3 & 4 \\ 9 & 2 & 0 \end{bmatrix}$.

Matrix Name:

Matrices are denoted conventionally by capital letters A, B, C... X, Y, Z etc of English Alphabets.

Row of a matrix:

In matrix, the entries presented in **horizontal** way are called rows.

$$A = \begin{bmatrix} a & b & c \\ \ell & m & b \end{bmatrix} \begin{matrix} \rightarrow R_1 \\ \rightarrow R_2 \end{matrix}$$

In above matrix A, R₁ and R₂ are two rows.

Columns of a matrix:

In matrix, the entries presented in **vertical** way are called columns.

$$A = \begin{bmatrix} 1 & 4 \\ 3 & 9 \end{bmatrix}$$

↓ ↓
C₁ C₂

In above matrix A, C₁ and C₂ are two columns.

Order of a Matrix:

The number of rows and columns in a matrix specifies its order.

The order of a matrix is denoted by m × n or m-by-n.

Here; “m” represented the number of rows and “n” represented the number of columns.

$$m - \text{by} - n$$

↓ ↓

No. of rows No. of columns .

If a matrix C has two rows and 3 columns. The order of matrix is 2-by-3.

$$C = \begin{bmatrix} 1 & 3 & 4 \\ 9 & 2 & 0 \end{bmatrix} \begin{matrix} \rightarrow R_1 \\ \rightarrow R_2 \end{matrix}$$

↓ ↓ ↓
C₁ C₂ C₃

The order of matrix C is 2-by-3.

Equal Matrices:

Let A and B be two matrices; if

- (i) The order of A = the order of B
 - (ii) Their corresponding entries are equal or same
- Then A and B are Equal matrices

Equal matrices are denoted by A = B

$$A = \begin{bmatrix} 2 & 4 \\ 9 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 2+2 \\ 10-1 & 1+1+1 \end{bmatrix}$$

Matrix A and B are equal because they have same order which is 2-by-2 and same corresponding elements so, A = B.

EXERCISE 1.1

Q.1: Find the order of the following matrices.

$$A = \begin{bmatrix} 2 & 3 \\ -5 & 6 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix},$$

$$C = [2 \quad 4], \quad D = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix},$$

$$E = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}, \quad F = [2]$$

$$G = \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 3 \\ 2 & 4 & 5 \end{bmatrix}, \quad H = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 0 & 6 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 3 \\ -5 & 6 \end{bmatrix}$$

Sol. $A = \begin{bmatrix} 2 & 3 \\ -5 & 6 \end{bmatrix}$

order of matrix = 2-by-2

$$B = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix}$$

Sol. $B = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix}$

order of matrix = 2-by-2

$$C = [2 \quad 4]$$

Sol. $C = [2 \quad 4]$

order of matrix = 1-by-2

$$D = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$$

Sol. $D = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$

order of matrix = 3-by-1

$$E = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}$$

Sol. $E = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}$

Order of matrix = 3-by-2.

$$F = [2]$$

Sol. $F = [2]$

Order of matrix = 1-by-1.

$$G = \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 3 \\ 2 & 4 & 5 \end{bmatrix}$$

Sol. $G = \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 3 \\ 2 & 4 & 5 \end{bmatrix}$

Order of matrix = 3-by-3.

$$H = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 0 & 6 \end{bmatrix}$$

Sol. $H = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 0 & 6 \end{bmatrix}$

Order of matrix = 2-by-3.

Q.2: Which of the following matrices are equal.

$A = [3], B = [3 \ 5], C = [5 \ -2],$

$D = [5 \ 3], E = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix},$

$F = \begin{bmatrix} 2 \\ 6 \end{bmatrix} \quad G = \begin{bmatrix} 3-1 \\ 3+3 \end{bmatrix},$

$H = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix}, I = [3 \ 3+2]$

$J = \begin{bmatrix} 2+2 & 2-2 \\ 2+4 & 2+0 \end{bmatrix},$

Sol. Since order of A and C same and corresponding elements are also same so $A = C$

$A = [3] \quad A = C$

$B = [3 \ 5] \quad B = I$

$C = [5 \ -2] \quad C = A$

$D = [5 \ 3] \quad \text{Not equal to any matrix}$

$E = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix} \quad E = J = H$

$F = \begin{bmatrix} 2 \\ 6 \end{bmatrix} \quad F = G$

$G = \begin{bmatrix} 3-1 \\ 3+3 \end{bmatrix} = \begin{bmatrix} 2 \\ 6 \end{bmatrix} \quad G = F$

$H = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix} \quad H = J = E$

$I = [3 \ 3+2] = [3 \ 5] \quad I = B$

$J = \begin{bmatrix} 2+2 & 2-2 \\ 2+4 & 2+0 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix} \quad J = H = E$

Q.3: Find the values of a, b, c and d which satisfy the matrix equation.

$\begin{bmatrix} a+c & a+2b \\ c-1 & 4d-6 \end{bmatrix} = \begin{bmatrix} 0 & -7 \\ 3 & 2d \end{bmatrix}$

Sol. $\begin{bmatrix} a+c & a+2b \\ c-1 & 4d-6 \end{bmatrix} = \begin{bmatrix} 0 & -7 \\ 3 & 2d \end{bmatrix}$

According to the definition of equal matrices.

$a + c = 0 \quad \dots\dots(i)$

$c - 1 = 3 \quad \dots\dots(ii)$

$a + 2b = -7 \quad \dots\dots(iii)$

$4d - 6 = 2d \quad \dots\dots(iv)$

By using the(ii) Put $c=4$ in (i) equation

$c - 1 = 3$

$c = 3+1$

$c = 4$

$a + c = 0$

$a + 4 = 0$

$a = 0 - 4$

$a = -4$

Put the value of $a = -4$ in (iii) equation

$a + 2b = -7$

$-4 + 2b = -7$

$2b = -7 + 4$

$2b = -3$

$b = -\frac{3}{2}$ or -1.5

By using (iv) equation

$4d - 6 = 2d$

$4d - 2d = 6$

$2d = 6$

$d = \frac{6}{2}$

$d = 3$

Hence the value of

$a = -4, b = -1.5, c = 4, d = 3$

TYPES OF MATRICES

(i) Row Matrix:

A matrix is called a row matrix if it has only one row.

e.g. $D = [1 \ 3 \ 4]$

D is a row matrix and its order is 1-by-3.

(ii) Column Matrix:

A matrix is called a column matrix if it has only one column

e.g; $E = \begin{bmatrix} 1 \\ 3 \\ 4 \\ 7 \end{bmatrix}$

E is a column matrix and its order is 4-by-1.

(iii) Rectangular Matrix:

A matrix A is called rectangular if, its number of rows is not equal to the number of its columns.

e.g; $A = \begin{bmatrix} 1 & 7 \\ 2 & 4 \\ 3 & 9 \end{bmatrix} \quad B = \begin{bmatrix} a & b & c \\ d & c & f \end{bmatrix}$

Order of A = 3-by-2

Order of B = 2-by-3

(iv) Square Matrix:

A matrix is called a square matrix if its number of rows is equal to its number of columns.

e.g; $C = [9] \quad D = \begin{bmatrix} 2 & 4 \\ 9 & 7 \end{bmatrix} \quad E = \begin{bmatrix} 1 & 7 & 0 \\ 3 & 9 & 3 \\ 5 & 11 & 2 \end{bmatrix}$

order 1-by-1 order 2-by-2 order 3-by-3

(v) Null or Zero Matrix:

A matrix is called a null or zero matrix if each of its entries/elements are zero (0).

e.g; $O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, O = [0 \ 0 \ 0], O = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

order 2-by-2 order 1-by-3 order 3-by-1

It is represented by O.

(vi) Transpose of a Matrix:

A matrix obtained by interchanging the row of matrix into the columns of that matrix.

OR

A matrix obtained by changing the columns into rows of a matrix.

If A is a matrix then transpose is denoted by A^t .

$$\text{e.g; } A = \begin{bmatrix} 0 & 4 \\ 1 & 5 \\ 2 & 7 \end{bmatrix} \quad \text{then } A^t = \begin{bmatrix} 0 & 1 & 2 \\ 4 & 5 & 7 \end{bmatrix}$$

order = 3-by-2 then transpose order = 2-by-3

(vii) Negative of a Matrix:

Let A be a matrix. Then its negative is obtained by changing the signs of all the elements of A, i.e.

$$\text{if } A = \begin{bmatrix} 2 & 9 \\ 4 & -3 \end{bmatrix}, \text{ then } -A = \begin{bmatrix} -2 & -9 \\ -4 & 3 \end{bmatrix}$$

(viii) Symmetric Matrix:

A square matrix is symmetric if it is equal to its transpose.

i.e; matrix A is symmetric if $A^t = A$.

$$A = \begin{bmatrix} a & b \\ b & a \end{bmatrix}, \quad A^t = \begin{bmatrix} a & b \\ b & a \end{bmatrix}$$

$$A^t = A$$

So, A is symmetric matrix.

(ix) Skew-Symmetric Matrix:

A square matrix A is said to be skew-symmetric if $A^t = -A$.

$$A = \begin{bmatrix} 0 & 2 & 3 \\ -2 & 0 & 1 \\ -3 & -1 & 0 \end{bmatrix}$$

$$\text{then } A^t = \begin{bmatrix} 0 & -2 & -3 \\ 2 & 0 & -1 \\ 3 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -2 & -3 \\ -(-2) & 0 & -1 \\ -(-3) & -(-1) & 0 \end{bmatrix}$$

$$= - \begin{bmatrix} 0 & 2 & 3 \\ -2 & 0 & 1 \\ -3 & -1 & 0 \end{bmatrix} = -A$$

Since $A^t = -A$, therefore A is skew-symmetric matrix.

(x) Diagonal Matrix:

A square matrix A is called a diagonal matrix if each element is zero except diagonal elements.

$$\text{e.g; } A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

(xi) Scalar Matrix:

A diagonal matrix is called a scalar matrix, if all the diagonal elements are same and non-zero.

$$A = \begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix} \quad k \neq 0, 1 \quad C = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 9 \end{bmatrix}$$

is a scalar matrix of order 3-by-3.

(xii) Identity Matrix:

A diagonal matrix is called identity (unit) matrix if all diagonal elements are 1 and it is denoted by I.

$$\text{e.g; } I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad I = [1]$$

are all unit matrices,

Remember:

Note: (i) The scalar matrix and identity matrix are diagonal matrix.

(ii) Every diagonal matrix is not a scalar or identity matrix.

EXERCISE 1.2

Q.1: From the following matrices, identify unit matrices row matrices, column matrices and null matrices.

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad B = [2 \quad 3 \quad 4],$$

$$C = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$E = [0], \quad F = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$$

Sol. $A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ is a null matrix.

$B = [2 \quad 3 \quad 4]$ is a row matrix.

$C = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$ is a column matrix.

$D = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ is a unit matrix.

$E = [0]$ is a null matrix.

$F = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$ is a column matrix.

Q.2: From the following matrices identify:

- (a) Square matrices
 (b) Rectangular matrices
 (c) Row matrices
 (d) Column matrices
 (d) Identity matrices
 (f) Null Matrices

(i) $\begin{bmatrix} -8 & 2 & 7 \\ 12 & 0 & 4 \end{bmatrix}$ (ii) $\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$ (iii) $\begin{bmatrix} 6 & -4 \\ 3 & -2 \end{bmatrix}$

(iv) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (v) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$ (vi) $[3 \ 10 \ -1]$

(vii) $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ (viii) $\begin{bmatrix} 1 & 2 & 3 \\ -1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (ix) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$

Sol. (i) $\begin{bmatrix} -8 & 2 & 7 \\ 12 & 0 & 4 \end{bmatrix}$ Rectangular matrix

(ii) $\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$ Column matrix, rectangular matrix

(iii) $\begin{bmatrix} 6 & -4 \\ 3 & -2 \end{bmatrix}$ Square matrix

(iv) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ Identity matrix, square matrix

(v) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$ Rectangular matrix

(vi) $[3 \ 10 \ -1]$ Row matrix, rectangular matrix

(vii) $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ Column matrix, rectangular matrix

(viii) $\begin{bmatrix} 1 & 2 & 3 \\ -1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ Square matrix

(ix) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ is a null matrix, rectangular matrix

Q.3: From the following matrices. Identify diagonal, scalar and unit (identity) matrices.

$A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

$D = \begin{bmatrix} 3 & 0 \\ 0 & 0 \end{bmatrix}$, $E = \begin{bmatrix} 5-3 & 0 \\ 0 & 1+1 \end{bmatrix}$

Sol. $A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}$ Scalar matrix, diagonal matrix

$B = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix}$ Diagonal matrix

$C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ Identity matrix, diagonal matrix

$D = \begin{bmatrix} 3 & 0 \\ 0 & 0 \end{bmatrix}$ Diagonal matrix

$E = \begin{bmatrix} 5-3 & 0 \\ 0 & 1+1 \end{bmatrix}$ $E = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ Scalar matrix, diagonal matrix

Q.4: Find negative of matrices A, B, C, D and E when

$A = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix}$, $C = \begin{bmatrix} 2 & 6 \\ 3 & 2 \end{bmatrix}$,

$D = \begin{bmatrix} -3 & 2 \\ -4 & 5 \end{bmatrix}$, $E = \begin{bmatrix} 1 & -5 \\ 2 & 3 \end{bmatrix}$

$A = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$

Sol. $A = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \Rightarrow -A = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$

$B = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix}$

Sol. $-B = \begin{bmatrix} -3 & 1 \\ -2 & -1 \end{bmatrix}$

$C = \begin{bmatrix} 2 & 6 \\ 3 & 2 \end{bmatrix}$

Sol. $-C = \begin{bmatrix} -2 & -6 \\ -3 & -2 \end{bmatrix}$

$D = \begin{bmatrix} -3 & 2 \\ -4 & 5 \end{bmatrix}$

Sol. $-D = \begin{bmatrix} 3 & -2 \\ 4 & -5 \end{bmatrix}$

$E = \begin{bmatrix} 1 & -5 \\ 2 & 3 \end{bmatrix}$

Sol. $-E = \begin{bmatrix} -1 & 5 \\ -2 & -3 \end{bmatrix}$

Q.5: Find the transpose of each of the following matrices.

$A = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}$, $B = [5 \ 1 \ -6]$, $C = \begin{bmatrix} 1 & 2 \\ 2 & -1 \\ 3 & 0 \end{bmatrix}$,

$D = \begin{bmatrix} 2 & 3 \\ 0 & 5 \end{bmatrix}$, $E = \begin{bmatrix} 2 & 3 \\ -4 & 5 \end{bmatrix}$, $F = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

$$\text{Sol. } A = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 0 & 1 & -2 \end{bmatrix}'$$

$$A^t = [0 \quad 1 \quad -2]$$

$$\text{Sol. } B = [5 \quad 1 \quad -6]$$

$$B^t = [5 \quad 1 \quad -6]^t$$

$$B^t = \begin{bmatrix} 5 \\ 1 \\ -6 \end{bmatrix}$$

$$\text{Sol. } C = \begin{bmatrix} 1 & 2 \\ 2 & -1 \\ 3 & 0 \end{bmatrix}$$

$$C^t = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 0 \end{bmatrix}'$$

$$C^t = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 0 \end{bmatrix}$$

$$\text{Sol. } D = \begin{bmatrix} 2 & 3 \\ 0 & 5 \end{bmatrix}$$

$$D^t = \begin{bmatrix} 2 & 3 \\ 0 & 5 \end{bmatrix}'$$

$$D^t = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix}$$

$$\text{Sol. } E = \begin{bmatrix} 2 & 3 \\ -4 & 5 \end{bmatrix}$$

$$E^t = \begin{bmatrix} 2 & 3 \\ -4 & 5 \end{bmatrix}'$$

$$E^t = \begin{bmatrix} 2 & -4 \\ 3 & 5 \end{bmatrix}$$

$$\text{Sol. } F = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$F^t = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}'$$

$$F^t = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

$$\text{Q.6: Verify that if } A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$\text{then (i) } (A^t)^t = A \quad \text{(ii) } (B^t)^t = B$$

$$\text{(i) } (A^t)^t = A$$

$$\text{Sol. } A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}'$$

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$(A^t)^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}'$$

$$(A^t)^t = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = A$$

$$\text{(ii) } (B^t)^t = B$$

$$\text{Sol. } B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$B^t = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}'$$

$$(B^t) = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$(B^t)^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}'$$

$$(B^t)^t = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} = B$$

Addition and Subtraction of Matrices:

1. Addition of Matrices:

Let A and B be any two matrices with real entries; Matrices A and B are conformable for addition, if they have same order. Addition is denoted by $A + B$ and is obtained by adding the entries of the matrix A to the corresponding entries of B.

$$A = \begin{bmatrix} 2 & 3 & 0 \\ 5 & 6 & 1 \\ 2 & 1 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 3 & -2 & 5 \\ -1 & 4 & 1 \\ 4 & 2 & -4 \end{bmatrix}$$

$$\begin{aligned} A+B &= \begin{bmatrix} 2 & 3 & 0 \\ 5 & 6 & 1 \\ 2 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 3 & -2 & 5 \\ -1 & 4 & 1 \\ 4 & 2 & -4 \end{bmatrix} \\ &= \begin{bmatrix} 2+3 & 3-2 & 0+5 \\ 5-1 & 6+4 & 1+1 \\ 2+4 & 1+2 & 3-4 \end{bmatrix} = \begin{bmatrix} 5 & 1 & 5 \\ 4 & 10 & 2 \\ 6 & 3 & -1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \text{Or } B+A &= \begin{bmatrix} 3 & -2 & 5 \\ -1 & 4 & 1 \\ 4 & 2 & -4 \end{bmatrix} + \begin{bmatrix} 2 & 3 & 0 \\ 5 & 6 & 1 \\ 2 & 1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 3+2 & -2+3 & 5+0 \\ -1+5 & 4+6 & 1+1 \\ 4+2 & 2+1 & -4+3 \end{bmatrix} = \begin{bmatrix} 5 & 1 & 5 \\ 4 & 10 & 2 \\ 6 & 3 & -1 \end{bmatrix} \end{aligned}$$

2. Subtraction of Matrices:

Let A and B any two matrices. Matrices A and B are conformable for subtraction, if they have same order represented by A - B and is obtained by subtracting the entries of the matrix B to the corresponding entries of A.

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 0 \end{bmatrix} \text{ and } B = \begin{bmatrix} 0 & 2 & 2 \\ -1 & 4 & 3 \end{bmatrix} \text{ are}$$

conformable for subtraction.

$$\begin{aligned} \text{i.e. } A-B &= \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 0 \end{bmatrix} - \begin{bmatrix} 0 & 2 & 2 \\ -1 & 4 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 2-0 & 3-2 & 4-2 \\ 1-(-1) & 5-4 & 0-3 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 2 \\ 2 & 1 & -3 \end{bmatrix} \end{aligned}$$

Some

3. Commutative and Associative Law of Addition of Matrices:

$A + B = B + A$ Commutative w.r.t +

$A+(B+C) = (A+B)+C$ Associative w.r.t +

4. Additive Identity of a Matrix:

If A and B are two matrices of same order and

$$A + B = A \quad \text{or} \quad B + A = A$$

then matrix B is called additive identity of matrix A.

For any matrix A and zero matrix O of same order, O is called additive identity of A as

$$A + O = A = O + A$$

5. Additive Inverse of a Matrix:

If A and B are two matrices of same order and

$$A + B = 0 = B + A$$

Then A and B are called additive inverse of each other.

Note: Additive inverse of any matrix A is $-A$ obtained by changing their signs of each element.

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad -A = \begin{bmatrix} -a & -b \\ -c & -d \end{bmatrix}$$

EXERCISE 1.3

Q.1. Which of the following matrices are conformable for addition?

$$A = \begin{bmatrix} 2 & 1 \\ -1 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 3 \\ 1 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 0 \\ 2 & -1 \\ 1 & -2 \end{bmatrix},$$

$$D = \begin{bmatrix} 2+1 \\ 3 \end{bmatrix}, \quad E = \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}, \quad F = \begin{bmatrix} 3 & 2 \\ 1+1 & -4 \\ 3+2 & 2+1 \end{bmatrix}$$

$$\text{Sol. } A = \begin{bmatrix} 2 & 1 \\ -1 & 3 \end{bmatrix} \quad 2 - \text{by} - 2$$

$$B = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \quad 2 - \text{by} - 1$$

$$C = \begin{bmatrix} 1 & 0 \\ 2 & -1 \\ 1 & -2 \end{bmatrix} \quad 3 - \text{by} - 2$$

$$D = \begin{bmatrix} 2+1 \\ 3 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix} \quad 2 - \text{by} 1$$

$$E = \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix} \quad 2 - \text{by} - 2$$

$$F = \begin{bmatrix} 3 & 2 \\ 1+1 & -4 \\ 3+2 & 2+1 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 2 & -4 \\ 5 & 3 \end{bmatrix} \quad 3 - \text{by} - 2$$

A and E are conformable for addition because their orders are same.

B and D are conformable for addition because their orders same.

C and F are conformable for addition because their orders are same.

Q.2: Find the additive inverse of following matrices.

$$A = \begin{bmatrix} 2 & 4 \\ -2 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 & -1 \\ 2 & -1 & 3 \\ 3 & -2 & 1 \end{bmatrix},$$

$$C = \begin{bmatrix} 4 \\ -2 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 \\ -3 & -2 \\ 2 & 1 \end{bmatrix},$$

$$E = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad F = \begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$$

$$\text{Sol. } A = \begin{bmatrix} 2 & 4 \\ -2 & 1 \end{bmatrix}$$

$$-A = \begin{bmatrix} -2 & -4 \\ 2 & -1 \end{bmatrix} \text{ is additive inverse of } A$$

$$\text{Sol. } \mathbf{B} = \begin{bmatrix} 1 & 0 & -1 \\ 2 & -1 & 3 \\ 3 & -2 & 1 \end{bmatrix}$$

$$-\mathbf{B} = \begin{bmatrix} -1 & 0 & 1 \\ -2 & 1 & -3 \\ -3 & 2 & -1 \end{bmatrix} \text{ is additive inverse of B}$$

$$\text{Sol. } \mathbf{C} = \begin{bmatrix} 4 \\ -2 \end{bmatrix}$$

$$-\mathbf{C} = \begin{bmatrix} -4 \\ 2 \end{bmatrix} \text{ is additive inverse of C}$$

$$\text{Sol. } \mathbf{D} = \begin{bmatrix} 1 & 0 \\ -3 & -2 \\ 2 & 1 \end{bmatrix} \text{ is additive inverse of D}$$

$$-\mathbf{D} = \begin{bmatrix} -1 & 0 \\ 3 & 2 \\ -2 & -1 \end{bmatrix}$$

$$\text{Sol. } \mathbf{E} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$-\mathbf{E} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \text{ is additive inverse of E}$$

$$\text{Sol. } \mathbf{F} = \begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$$

$$-\mathbf{F} = \begin{bmatrix} -\sqrt{3} & -1 \\ 1 & -\sqrt{2} \end{bmatrix} \text{ is additive inverse of F}$$

Q.3. If $\mathbf{A} = \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix}$, $\mathbf{B} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$

$\mathbf{C} = [1 \quad -1 \quad 2]$, $\mathbf{D} = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix}$ then

find, (i) $\mathbf{A} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ (ii) $\mathbf{B} + \begin{bmatrix} -2 \\ 3 \end{bmatrix}$

(iii) $\mathbf{C} + [-2 \quad 1 \quad 3]$ (iv) $\mathbf{D} + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$

(v) $2\mathbf{A}$ (vi) $(-1)\mathbf{B}$ (vii) $(-2)\mathbf{C}$
 (viii) $3\mathbf{D}$ (ix) $3\mathbf{C}$

(i) $\mathbf{A} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

$$\text{Sol. } = \mathbf{A} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -1+1 & 2+1 \\ 2+1 & 1+1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 3 \\ 3 & 2 \end{bmatrix}$$

(ii) $\mathbf{B} + \begin{bmatrix} -2 \\ 3 \end{bmatrix}$

$$\text{Sol. } = \mathbf{B} + \begin{bmatrix} -2 \\ 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1 \\ -1 \end{bmatrix} + \begin{bmatrix} -2 \\ 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1-2 \\ -1+3 \end{bmatrix}$$

$$= \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

(iii) $\mathbf{C} + [-2 \quad 1 \quad 3]$

$$\text{Sol. } = \mathbf{C} + [-2 \quad 1 \quad 3]$$

$$= [1 \quad -1 \quad 2] + [-2 \quad 1 \quad 3]$$

$$= [1-2 \quad -1+1 \quad 2+3]$$

$$= [-1 \quad 0 \quad 5]$$

(iv) $\mathbf{D} + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$

$$\text{Sol. } = \mathbf{D} + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix} + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+0 & 2+1 & 3+0 \\ -1+2 & 0+0 & 2+1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 3 & 3 \\ 1 & 0 & 3 \end{bmatrix}$$

(v) $2\mathbf{A}$

$$\text{Sol. } = 2\mathbf{A}$$

$$= 2 \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} (2)(-1) & (2)(2) \\ (2)(2) & (2)(1) \end{bmatrix}$$

$$2\mathbf{A} = \begin{bmatrix} -2 & 4 \\ 4 & 2 \end{bmatrix}$$

(vi) $(-1)\mathbf{B}$

$$\text{Sol. } = -1 \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$= \begin{bmatrix} (-1)(1) \\ (-1)(-1) \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

(vii) $(-2)\mathbf{C}$

$$\text{Sol. } = -2[1 \quad -1 \quad 2]$$

$$= [(-2)(1) \quad (-2)(-1) \quad (-2)(2)]$$

$$= [-2 \quad 2 \quad -4]$$

(viii) $3\mathbf{D}$

$$\text{Sol. } = 3 \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} (3)(1) & (3)(2) & (3)(3) \\ (3)(-1) & (3)(0) & (3)(2) \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 6 & 9 \\ -3 & 0 & 6 \end{bmatrix}$$

(ix) 3C

$$\begin{aligned}\text{Sol.} &= 3[1 \quad -1 \quad 2] \\ &= [(3)(1) \quad (3)(-1) \quad (3)(2)] \\ &= [3 \quad -3 \quad 6]\end{aligned}$$

Q.4: Perform the indicated operations and simplify the following:

Two matrices are said to conformable for addition and subtraction if their orders same.

$$(i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$\begin{aligned}\text{Sol.} &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1+0 & 0+2 \\ 0+3 & 1+0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1+1 & 2+1 \\ 3+1 & 1+0 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}\end{aligned}$$

$$(ii) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$\begin{aligned}\text{Sol.} &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1+0 & 0+2 \\ 0+3 & 1+0 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1-1 & 2-1 \\ 3-1 & 1-0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix}\end{aligned}$$

$$(iii) [2 \quad 3 \quad 1] + ([1 \quad 0 \quad 2] - [2 \quad 2 \quad 2])$$

$$\begin{aligned}\text{Sol.} &= [2 \quad 3 \quad 1] + ([1 \quad 0 \quad 2] - [2 \quad 2 \quad 2]) \\ &= [2 \quad 3 \quad 1] + ([1-2 \quad 0-2 \quad 2-2]) \\ &= [2 \quad 3 \quad 1] + [-1 \quad -2 \quad 0] \\ &= [2-1 \quad 3-2 \quad 1+0] \\ &= [1 \quad 1 \quad 1]\end{aligned}$$

$$(iv) \begin{bmatrix} 1 & 2 & 3 \\ -1 & -1 & -1 \\ 0 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \end{bmatrix}$$

$$\begin{aligned}\text{Sol.} &= \begin{bmatrix} 1 & 2 & 3 \\ -1 & -1 & -1 \\ 0 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 1+1 & 2+1 & 3+1 \\ -1+2 & -1+2 & -1+2 \\ 0+3 & 1+3 & 2+3 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 3 & 4 \\ 1 & 1 & 1 \\ 3 & 4 & 5 \end{bmatrix}\end{aligned}$$

$$(v) \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & -2 \\ -2 & -1 & 0 \\ 0 & 2 & -1 \end{bmatrix}$$

$$\begin{aligned}\text{Sol.} &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & -2 \\ -2 & -1 & 0 \\ 0 & 2 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 1+1 & 2+0 & 3-2 \\ 2-2 & 3-1 & 1+0 \\ 3+0 & 1+2 & 2-1 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 2 & 1 \\ 0 & 2 & 1 \\ 3 & 3 & 1 \end{bmatrix}\end{aligned}$$

$$(vi) \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2+1 & 2+1+1 \\ 0+1+1 & 1+0+1 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 4 \\ 2 & 2 \end{bmatrix}$$

Q.5: For the matrices $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$,

$$B = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \text{ and } C = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

verify the following rules:

(i) $A + C = C + A$

Sol. L.H.S.

$$A + C = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1-1 & 2+0 & 3+0 \\ 2+0 & 3-2 & 1+3 \\ 1+1 & -1+1 & 0+2 \end{bmatrix} = \begin{bmatrix} 0 & 2 & 3 \\ 2 & 1 & 4 \\ 2 & 0 & 2 \end{bmatrix}$$

R.H.S.

$$C + A = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$C + A = \begin{bmatrix} -1+1 & 0+2 & 0+3 \\ 0+2 & -2+3 & 3+1 \\ 1+1 & 1-1 & 2+0 \end{bmatrix}$$

$$C + A = \begin{bmatrix} 0 & 2 & 3 \\ 2 & 1 & 4 \\ 2 & 0 & 2 \end{bmatrix}$$

Hence proved

L.H.S = R.H.S

(ii) $A + B = B + A$

Sol. L.H.S

$$A + B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$A + B = \begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3-2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix}$$

$$A + B = \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

R.H.S

$$B + A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$B + A = \begin{bmatrix} 1+1 & -1+2 & 1+3 \\ 2+2 & -2+3 & 2+1 \\ 3+1 & 1-1 & 3+0 \end{bmatrix}$$

$$B + A = \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

Hence proved L.H.S = R.H.S

(iii) $B + C = C + B$

Sol. L.H.S

$$B + C = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$B + C = \begin{bmatrix} 1-1 & -1+0 & 1+0 \\ 2+0 & -2-2 & 2+3 \\ 3+1 & 1+1 & 3+2 \end{bmatrix}$$

$$B + C = \begin{bmatrix} 0 & -1 & 1 \\ 2 & -4 & 5 \\ 4 & 2 & 5 \end{bmatrix}$$

R.H.S

$$C + B = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$C + B = \begin{bmatrix} -1+1 & 0-1 & 0+1 \\ 0+2 & -2-2 & 3+2 \\ 1+3 & 1+1 & 2+3 \end{bmatrix}$$

$$C + B = \begin{bmatrix} 0 & -1 & 1 \\ 2 & -4 & 5 \\ 4 & 2 & 5 \end{bmatrix}$$

Hence proved L.H.S = R.H.S

(iv) $A + (B + A) = 2A + B$

Sol. L.H.S:

$$A + (B + A)$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} \right)$$

$$A + (B + A)$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1+1 & -1+2 & 1+3 \\ 2+2 & -2+3 & 2+1 \\ 3+1 & 1-1 & 3+0 \end{bmatrix} \right)$$

$$A + (B + A) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2 & 2+1 & 3+4 \\ 2+4 & 3+1 & 1+3 \\ 1+4 & -1+0 & 0+3 \end{bmatrix}$$

$$A + (B + A) = \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix}$$

R.H.S:

$$2A + B = 2 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + B$$

$$= \begin{bmatrix} 2 \times 1 & 2 \times 2 & 2 \times 3 \\ 2 \times 2 & 2 \times 3 & 2 \times 1 \\ 2 \times 1 & 2 \times -1 & 2 \times 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} 2 & 4 & 6 \\ 4 & 6 & 2 \\ 2 & -2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} 2+1 & 4-1 & 6+1 \\ 4+2 & 6-2 & 2+2 \\ 2+3 & -2+1 & 0+3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix}$$

Hence proved L.H.S = R.H.S

(v) $(C - B) + A = C + (A - B)$

Sol. L.H.S:

$$(C - B) + A$$

$$= \left(\begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$(C - B) + A = \left(\begin{bmatrix} -1-1 & 0+1 & 0-1 \\ 0-2 & -2+2 & 3-2 \\ 1-3 & 1-1 & 2-3 \end{bmatrix} \right) + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$(C - B) + A = \begin{bmatrix} -2 & 1 & -1 \\ -2 & 0 & 1 \\ -2 & 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$(C - B) + A = \begin{bmatrix} -2+1 & 1+2 & -1+3 \\ -2+2 & 0+3 & 1+1 \\ -2+1 & 0-1 & -1+0 \end{bmatrix} = \begin{bmatrix} -1 & 3 & 2 \\ 0 & 3 & 2 \\ -1 & -1 & -1 \end{bmatrix}$$

R.H.S: C + (A - B)

C + (A - B)

$$= \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right)$$

$$C + (A - B) = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1-1 & 2+1 & 3-1 \\ 2-2 & 3+2 & 1-2 \\ 1-3 & -1-1 & 0-3 \end{bmatrix}$$

$$C + (A - B) = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 0 & 3 & 2 \\ 0 & 5 & -1 \\ -2 & -2 & -3 \end{bmatrix}$$

$$C + (A - B) = \begin{bmatrix} -1+0 & 0+3 & 0+2 \\ 0+0 & -2+5 & 3-1 \\ 1-2 & 1-2 & 2-3 \end{bmatrix}$$

$$C + (A - B) = \begin{bmatrix} -1 & 3 & 2 \\ 0 & 3 & 2 \\ -1 & -1 & -1 \end{bmatrix}$$

Hence proved.

L.H.S = R.H.S

(vi) $2A + B = A + (A + B)$

Sol: L.H.S:

$$2A + B = 2 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} (2)(1) & (2)(2) & (2)(3) \\ (2)(2) & (2)(3) & (2)(1) \\ (2)(1) & (2)(-1) & (2)(0) \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} 2 & 4 & 6 \\ 4 & 6 & 2 \\ 2 & -2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} 2+1 & 4-1 & 6+1 \\ 4+2 & 6-2 & 2+2 \\ 2+3 & -2+1 & 0+3 \end{bmatrix}$$

$$2A + B = \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix}$$

R.H.S:

A + (A + B)

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right)$$

A + (A + B)

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3-2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix}$$

$$A + (A + B) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

$$A + (A + B) = \begin{bmatrix} 1+2 & 2+1 & 3+4 \\ 2+4 & 3+1 & 1+3 \\ 1+4 & -1+0 & 0+3 \end{bmatrix}$$

$$A + (A + B) = \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix}$$

Hence proved. L.H.S = R.H.S

(vii) $(C - B) - A = (C - A) - B$

Sol. L.H.S:

(C - B) - A

$$= \left(\begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

(C - B) - A

$$= \left(\begin{bmatrix} -1-1 & 0+1 & 0-1 \\ 0-2 & -2+2 & 3-2 \\ 1-3 & 1-1 & 2-3 \end{bmatrix} \right) - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

(C - B) - A

$$= \begin{bmatrix} -2 & 1 & -1 \\ -2 & 0 & 1 \\ -2 & 0 & -1 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$(C - B) - A = \begin{bmatrix} -2-1 & 1-2 & -1-3 \\ -2-2 & 0-3 & 1-1 \\ -2-1 & 0+1 & -1-0 \end{bmatrix}$$

$$(C - B) - A = \begin{bmatrix} -3 & -1 & -4 \\ -4 & -3 & 0 \\ -3 & 1 & -1 \end{bmatrix}$$

R.H.S:

$$= \left(\begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} \right) - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \left(\begin{bmatrix} -1-1 & 0-2 & 0-3 \\ 0-2 & -2-3 & 3-1 \\ 1-1 & 1+1 & 2-0 \end{bmatrix} \right) - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$(C - A) - B = \begin{bmatrix} -2 & -2 & -3 \\ -2 & -5 & 2 \\ 0 & 2 & 2 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$(C - A) - B = \begin{bmatrix} -2-1 & -2+1 & -3-1 \\ -2-2 & -5+2 & 2-2 \\ 0-3 & 2-1 & 2-3 \end{bmatrix}$$

$$(C - A) - B = \begin{bmatrix} -3 & -1 & -4 \\ -4 & -3 & 0 \\ -3 & 1 & -1 \end{bmatrix}$$

Hence proved

L.H.S = R.H.S

(viii) $(A + B) + C = A + (B + C)$

Sol. L.H.S:

$$(A + B) + C = \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$(A + B) + C = \left(\begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3-2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix} \right) + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$(A + B) + C = \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$(A + B) + C = \begin{bmatrix} 2-1 & 1+0 & 4+0 \\ 4+0 & 1-2 & 3+3 \\ 4+1 & 0+1 & 3+2 \end{bmatrix}$$

$$(A + B) + C = \begin{bmatrix} 1 & 1 & 4 \\ 4 & -1 & 6 \\ 5 & 1 & 5 \end{bmatrix}$$

R.H.S:

$A + (B + C)$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \right)$$

$$A + (B + C) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1-1 & -1+0 & 1+0 \\ 2+0 & -2-2 & 2+3 \\ 3+1 & 1+1 & 3+2 \end{bmatrix} \right)$$

$$A + (B + C) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 0 & -1 & 1 \\ 2 & -4 & 5 \\ 4 & 2 & 5 \end{bmatrix}$$

$$A + (B + C) = \begin{bmatrix} 1+0 & 2-1 & 3+1 \\ 2+2 & 3-4 & 1+5 \\ 1+4 & -1+2 & 0+5 \end{bmatrix}$$

$$A + (B + C) = \begin{bmatrix} 1 & 1 & 4 \\ 4 & -1 & 6 \\ 5 & 1 & 5 \end{bmatrix}$$

Hence proved.

L.H.S = R.H.S

(ix) $A + (B - C) = (A - C) + B$

Sol. L.H.S:

$$A + (B - C) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} - \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \right)$$

$$A + (B - C) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1+1 & -1-0 & 1-0 \\ 2-0 & -2+2 & 2-3 \\ 3-1 & 1-1 & 3-2 \end{bmatrix} \right)$$

$$A + (B - C) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & -1 & 1 \\ 2 & 0 & -1 \\ 2 & 0 & 1 \end{bmatrix}$$

$$A + (B - C) = \begin{bmatrix} 1+2 & 2-1 & 3+1 \\ 2+2 & 3+0 & 1-1 \\ 1+2 & -1+0 & 0+1 \end{bmatrix}$$

$$A + (B - C) = \begin{bmatrix} 3 & 1 & 4 \\ 4 & 3 & 0 \\ 3 & -1 & 1 \end{bmatrix}$$

R.H.S:

$(A - C) + B$

$$= \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} - \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \right) + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$(A - C) + B = \left(\begin{bmatrix} 1+1 & 2-0 & 3-0 \\ 2-0 & 3+2 & 1-3 \\ 1-1 & -1-1 & 0-2 \end{bmatrix} \right) + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$(A-C)+B = \begin{bmatrix} 2 & 2 & 3 \\ 2 & 5 & -2 \\ 0 & -2 & -2 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$(A-C)+B = \begin{bmatrix} 2+1 & 2-1 & 3+1 \\ 2+2 & 5-2 & -2+2 \\ 0+3 & -2+1 & -2+3 \end{bmatrix}$$

$$(A-C)+B = \begin{bmatrix} 3 & 1 & 4 \\ 4 & 3 & 0 \\ 3 & -1 & 1 \end{bmatrix}$$

Hence proved.

L.H.S = R.H.S

(x) $2A + 2B = 2(A + B)$

Sol. L.H.S:

$$2A + 2B = 2 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + 2 \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$2A + 2B = \begin{bmatrix} (2)(1) & (2)(2) & (2)(3) \\ (2)(2) & (2)(3) & (2)(1) \\ (2)(1) & (2)(-1) & (2)(0) \end{bmatrix} + \begin{bmatrix} (2)(1) & (2)(-1) & (2)(1) \\ (2)(2) & (2)(-2) & (2)(2) \\ (2)(3) & (2)(1) & (2)(3) \end{bmatrix}$$

$$2A + 2B = \begin{bmatrix} 2 & 4 & 6 \\ 4 & 6 & 2 \\ 2 & -2 & 0 \end{bmatrix} + \begin{bmatrix} 2 & -2 & 2 \\ 4 & -4 & 4 \\ 6 & 2 & 6 \end{bmatrix}$$

$$2A + 2B = \begin{bmatrix} 2+2 & 4-2 & 6+2 \\ 4+4 & 6-4 & 2+4 \\ 2+6 & -2+2 & 0+6 \end{bmatrix}$$

$$2A + 2B = \begin{bmatrix} 4 & 2 & 8 \\ 8 & 2 & 6 \\ 8 & 0 & 6 \end{bmatrix}$$

R.H.S:

$$2(A + B) = 2 \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right)$$

$$2(A + B) = 2 \left(\begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3-2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix} \right)$$

$$2(A + B) = 2 \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

$$2(A + B) = \begin{bmatrix} 2(2) & 1(2) & 4(2) \\ 4(2) & 1(2) & 3(2) \\ 4(2) & 0(2) & 3(2) \end{bmatrix}$$

$$2(A + B) = \begin{bmatrix} 4 & 2 & 8 \\ 8 & 2 & 6 \\ 8 & 0 & 6 \end{bmatrix}$$

Hence proved. L.H.S = R.H.S

Q.6: If $A = \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 7 \\ -3 & 8 \end{bmatrix}$ find.

(i) $3A - 2B$ (ii) $2A^t - 3B^t$

(i) $3A - 2B$

Sol.

$$3A - 2B = 3 \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix} - 2 \begin{bmatrix} 0 & 7 \\ -3 & 8 \end{bmatrix}$$

$$3A - 2B = \begin{bmatrix} 3(1) & 3(-2) \\ 3(3) & 3(4) \end{bmatrix} - \begin{bmatrix} 2(0) & 2(7) \\ 2(-3) & 2(8) \end{bmatrix}$$

$$3A - 2B = \begin{bmatrix} 3 & -6 \\ 9 & 12 \end{bmatrix} - \begin{bmatrix} 0 & 14 \\ -6 & 16 \end{bmatrix}$$

$$3A - 2B = \begin{bmatrix} 3-0 & -6-14 \\ 9+6 & 12-16 \end{bmatrix} = \begin{bmatrix} 3 & -20 \\ 15 & -4 \end{bmatrix}$$

(ii) $2A^t - 3B^t$

Sol. $A = \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix} \Rightarrow A^t = \begin{bmatrix} 1 & 3 \\ -2 & 4 \end{bmatrix}$

$$B = \begin{bmatrix} 0 & 7 \\ -3 & 8 \end{bmatrix} \Rightarrow B^t = \begin{bmatrix} 0 & -3 \\ 7 & 8 \end{bmatrix}$$

$$2A^t - 3B^t = 2 \begin{bmatrix} 1 & 3 \\ -2 & 4 \end{bmatrix} - 3 \begin{bmatrix} 0 & -3 \\ 7 & 8 \end{bmatrix}$$

$$2A^t - 3B^t = \begin{bmatrix} 2 & 6 \\ -4 & 8 \end{bmatrix} - \begin{bmatrix} 0 & -9 \\ 21 & 24 \end{bmatrix}$$

$$2A^t - 3B^t = \begin{bmatrix} 2-0 & 6+9 \\ -4-21 & 8-24 \end{bmatrix}$$

$$2A^t - 3B^t = \begin{bmatrix} 2 & 15 \\ -25 & -16 \end{bmatrix}$$

Q.7. If $2 \begin{bmatrix} 2 & 4 \\ -3 & a \end{bmatrix} + 3 \begin{bmatrix} 1 & b \\ 8 & -4 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix}$ then find a and b.

Sol. $2 \begin{bmatrix} 2 & 4 \\ -3 & a \end{bmatrix} + 3 \begin{bmatrix} 1 & b \\ 8 & -4 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix}$

$$\begin{bmatrix} 4 & 8 \\ -6 & 2a \end{bmatrix} + \begin{bmatrix} 3 & 3b \\ 24 & -12 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 4+3 & 8+3b \\ -6+24 & 2a-12 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 7 & 8+3b \\ 18 & 2a-12 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix}$$

According to the definition of equal matrices

$$8 + 3b = 10 \quad 2a - 12 = 1 \dots\dots(ii)$$

$$3b = 10 - 8 \quad 2a = 1 + 12$$

$$3b = 2 \quad 2a = 13$$

$$b = \frac{2}{3} \quad a = \frac{13}{2}$$

Q.8: If $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$ then verify

that

$$(i) (A + B)^t = A^t + B^t$$

L.H.S

$$= (A + B)^t$$

Sol.

$$= (A+B)$$

$$(A + B) = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$(A + B) = \begin{bmatrix} 1+1 & 2+1 \\ 0+2 & 1+0 \end{bmatrix}$$

$$(A + B) = \begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix}$$

$$\text{Now } (A + B)^t = \begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix}^t$$

$$(A + B)^t = \begin{bmatrix} 2 & 2 \\ 3 & 1 \end{bmatrix}$$

R.H.S

$$\therefore A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}, B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$A^t + B^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 0+2 \\ 2+1 & 1+0 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 2 \\ 3 & 1 \end{bmatrix}$$

Hence proved. L.H.S = R.H.S

$$(ii) (A - B)^t = A^t - B^t$$

L.H.S

$$(A - B)^t$$

$$A - B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$A - B = \begin{bmatrix} 1-1 & 2-1 \\ 0-2 & 1-0 \end{bmatrix}$$

$$(A - B) = \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix}$$

$$(A - B)^t = \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix}^t$$

$$(A - B)^t = \begin{bmatrix} 0 & -2 \\ 1 & 1 \end{bmatrix}$$

R.H.S

$$= A^t - B^t$$

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}, B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$A^t - B^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$A^t - B^t = \begin{bmatrix} 1-1 & 0-2 \\ 2-1 & 1-0 \end{bmatrix}$$

$$A^t - B^t = \begin{bmatrix} 0 & -2 \\ 1 & 1 \end{bmatrix}$$

Hence proved L.H.S = R.H.S

$$(iii) A + A^t \text{ is symmetric } A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$\text{Sol. } A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$A + A^t = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$A + A^t = \begin{bmatrix} 1+1 & 2+0 \\ 0+2 & 1+1 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

$$(A + A^t)^t = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}^t$$

$$(A + A^t)^t = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

So proved according to the definition of symmetric matrix $(A + A^t) = (A + A^t)^t$

$$(iv) A - A^t \text{ is a skew symmetric } A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$\text{Sol. } A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$A - A^t = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$A - A^t = \begin{bmatrix} 1-1 & 2-0 \\ 0-2 & 1-1 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

$$(A - A^t)^t = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}^t$$

$$\text{Now } -(A - A^t) = - \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

$$(A - A^t)^t = \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix} = - \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} = -(A - A^t)^t$$

So proved according to the definition of skew symmetric matrix $(A - A^t) = -(A - A^t)^t$

$$(v) B + B^t \text{ is symmetric matrix } B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$\text{Sol. } B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$B + B^t = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$B + B^t = \begin{bmatrix} 1+1 & 1+2 \\ 2+1 & 0+0 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 3 & 0 \end{bmatrix}$$

$$(B + B^t)^t = \begin{bmatrix} 2 & 3 \\ 3 & 0 \end{bmatrix}^t$$

$$(B + B^t)^t = \begin{bmatrix} 2 & 3 \\ 3 & 0 \end{bmatrix} = -(B + B^t)$$

So according to the definition of symmetric matrix, $B + B^t$ is symmetric matrix.

(vi) $B - B^t$ is symmetric $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$

Sol. $B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$

$$B - B^t = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1-1 & 1-2 \\ 2-1 & 0-0 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

To prove that $(B - B^t)$ is skew symmetric matrix of $(B - B^t)$

$$(B - B^t)^t = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^t$$

$$(B - B^t)^t = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = -\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

So $B - B^t$ is skew symmetric matrix.

EXERCISE 1.4

Note: Two Matrices A and B are conformable for multiplication (giving product AB) if number of columns of 1st Matrix (i.e. A) equal to the number of rows of 2nd Matrix (i.e. B).

Q.1: Which of the following product of matrices is conformable for multiplication?

(i) $\begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -2 \\ 3 \end{bmatrix}$

No. of columns of 1st Matrix = 2

No. of rows of 2nd Matrix = 2

Sol. $2-by-2 \quad 2-by-1$

So it is possible

(ii) $\begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$

No. of columns of 1st Matrix = 2

No. of rows of 2nd Matrix = 2

Sol. $2-by-2 \quad 2-by-2$

So it is possible

(iii) $\begin{bmatrix} 1 \\ -1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$

No. of columns of 1st Matrix = 1

No. of rows of 2nd Matrix = 2

Sol. $2-by-1 \quad 2-by-2$

So it is impossible

(iv) $\begin{bmatrix} 1 & 2 \\ 0 & -1 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$

No. of columns of 1st Matrix = 2

No. of rows of 2nd Matrix = 2

Sol. $3-by-2 \quad 2-by-3$

So it is possible

(v) $\begin{bmatrix} 3 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ -2 & 3 \end{bmatrix}$

No. of columns of 1st Matrix = 3

No. of rows of 2nd Matrix = 3

Sol. $2-by-3 \quad 3-by-2$

So it is possible

Q.2. If $A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 6 \\ 5 \end{bmatrix}$, find (i) AB (ii) BA (if possible).

(i) AB

Sol. $A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 6 \\ 5 \end{bmatrix}$ product is possible.

So $AB = \begin{bmatrix} 3 & 0 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 6 \\ 5 \end{bmatrix}$

$$AB = \begin{bmatrix} (3)(6) + (0)(5) \\ (-1)(6) + (2)(5) \end{bmatrix}$$

$$AB = \begin{bmatrix} 18 + 0 \\ -6 + 10 \end{bmatrix} \Rightarrow AB = \begin{bmatrix} 18 \\ 4 \end{bmatrix}$$

(ii) BA

Sol. $A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 6 \\ 5 \end{bmatrix}$

BA is impossible. \because According to rule

No. of columns of 1st = 1

No. of rows of 2nd = 2

So Impossible

Q.3: Find the following product.

(i) $\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \end{bmatrix}$

Sol. $= \begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \end{bmatrix} \downarrow$

$$= [(1)(4) + (2)(0)]$$

$$= [4 + 0]$$

$$= [4]$$

(ii) $[1 \ 2] \begin{bmatrix} 5 \\ -4 \end{bmatrix}$

Sol. $= [1 \ 2] \begin{bmatrix} 5 \\ -4 \end{bmatrix} \downarrow$
 $= [(1)(5) + (2)(-4)]$
 $= [5 - 8]$
 $= [-3]$

(iii) $[-3 \ 0] \begin{bmatrix} 4 \\ 0 \end{bmatrix}$

Sol. $= [-3 \ 0] \begin{bmatrix} 4 \\ 0 \end{bmatrix} \downarrow$
 $= [(-3)(4) + (0)(0)]$
 $= [-12 + 0]$
 $= [-12]$

(iv) $[6 \ -0] \begin{bmatrix} 4 \\ 0 \end{bmatrix}$

Sol. $= [6 \ -0] \begin{bmatrix} 4 \\ 0 \end{bmatrix} \downarrow$
 $= [(6)(4) + (-0)(0)]$
 $= [24 - 0]$
 $= [24]$

(v) $\begin{bmatrix} 1 & 2 \\ -3 & 0 \\ 6 & -1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 0 & -4 \end{bmatrix}$

Sol. $= \begin{bmatrix} 1 & 2 \\ -3 & 0 \\ 6 & -1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 0 & -4 \end{bmatrix} \downarrow$
 $= \begin{bmatrix} (1)(4)+(2)(0) & (1)(5)+(2)(-4) \\ (-3)(4)+(0)(0) & (-3)(5)+(0)(-4) \\ (6)(4)+(-1)(0) & (6)(5)+(-1)(-4) \end{bmatrix}$
 $= \begin{bmatrix} 4+0 & 5-8 \\ -12+0 & -15-0 \\ 24-0 & 30+4 \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ -12 & -15 \\ 24 & 34 \end{bmatrix}$

Q.4: Multiply the following matrices.

(a) $\begin{bmatrix} 2 & 3 \\ 1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 3 & 0 \end{bmatrix}$

Sol. $= \begin{bmatrix} 2 & 3 \\ 1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 3 & 0 \end{bmatrix}$
 $= \begin{bmatrix} 2 \ 2 + 3 \ 3 & 2 \ -1 + 3 \ 0 \\ 1 \ 2 + 1 \ 3 & 1 \ -1 + 1 \ 0 \\ 0 \ 2 + -2 \ 3 & 0 \ -1 + -2 \ 0 \end{bmatrix}$

$$= \begin{bmatrix} 4+9 & -2+0 \\ 2+3 & -1+0 \\ 0-6 & 0+0 \end{bmatrix}$$

$$= \begin{bmatrix} 13 & -2 \\ 5 & -1 \\ -6 & 0 \end{bmatrix}$$

4. (b) $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix}$

Sol. $= \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix} \downarrow$
 $= \begin{bmatrix} (1)(1) + (2)(3) + (3)(-1) & (1)(2) + (2)(4) + (3)(1) \\ (4)(1) + (5)(3) + (6)(-1) & (4)(2) + (5)(4) + (6)(1) \end{bmatrix}$
 $= \begin{bmatrix} 1+6-3 & 2+8+3 \\ 4+15-6 & 8+20+6 \end{bmatrix}$
 $= \begin{bmatrix} 4 & 13 \\ 13 & 34 \end{bmatrix}$

4.(c) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$

Sol. $= \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \downarrow$
 $= \begin{bmatrix} (1)(1) + (2)(4) & (1)(2) + (2)(5) & (1)(3) + (2)(6) \\ (3)(1) + (4)(4) & (3)(2) + (4)(5) & (3)(3) + (4)(6) \\ (-1)(1) + (1)(4) & (-1)(2) + (1)(5) & (-1)(3) + (1)(1) \end{bmatrix}$
 $= \begin{bmatrix} 1+8 & 2+10 & 3+12 \\ 3+16 & 6+20 & 9+24 \\ -1+4 & -2+5 & -3+6 \end{bmatrix}$
 $= \begin{bmatrix} 9 & 12 & 15 \\ 19 & 26 & 33 \\ 3 & 3 & 3 \end{bmatrix}$

4. (d) $\begin{bmatrix} 8 & 5 \\ 6 & 4 \end{bmatrix} \begin{bmatrix} 2 & -5 \\ -4 & 4 \end{bmatrix}$

Sol. $= \begin{bmatrix} 8 & 5 \\ 6 & 4 \end{bmatrix} \begin{bmatrix} 2 & -5 \\ -4 & 4 \end{bmatrix} \downarrow$
 $= \begin{bmatrix} (8)(2) + (5)(-4) & (8)(\frac{-5}{2}) + (5)(4) \\ (6)(2) + (4)(-4) & (6)(\frac{-5}{2}) + (4)(4) \end{bmatrix}$

$$= \begin{bmatrix} 16-20 & -20+20 \\ 12-16 & -15+16 \end{bmatrix}$$

$$= \begin{bmatrix} -4 & 0 \\ -4 & 1 \end{bmatrix}$$

$$4. (e) \begin{bmatrix} -1 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{Sol.} = \begin{bmatrix} \overrightarrow{-1} & \overrightarrow{2} \\ \overrightarrow{1} & \overrightarrow{3} \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \downarrow$$

$$= \begin{bmatrix} (-1)(0)+(2)(0) & (-1)(0)+(2)(0) \\ (1)(0)+(3)(0) & (1)(0)+(3)(0) \end{bmatrix}$$

$$= \begin{bmatrix} 0+0 & 0+0 \\ 0+0 & 0+0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Q.5: Let $A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$ verify that:

(i) $AB = BA$ It is called commutative law w.r.t multiplication.

Note: In Matrices this law does not hold generally. i.e $AB \neq BA$

Sol. $AB = BA$

L.H.S

$$AB = \begin{bmatrix} \overrightarrow{-1} & \overrightarrow{3} \\ \overrightarrow{2} & \overrightarrow{0} \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} \downarrow$$

$$AB = \begin{bmatrix} (-1)(1)+(3)(-3) & (-1)(2)+(3)(-5) \\ (2)(1)+(0)(-3) & (2)(2)+(0)(-5) \end{bmatrix}$$

$$AB = \begin{bmatrix} -1-9 & -2-15 \\ 2-0 & 4-0 \end{bmatrix}$$

$$AB = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}$$

R.H.S

$$BA = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$$

$$BA = \begin{bmatrix} (1)(-1)+(2)(2) & (1)(3)+(2)(0) \\ (-3)(-1)+(-5)(2) & (-3)(3)+(-5)(0) \end{bmatrix}$$

$$BA = \begin{bmatrix} -1+4 & 3+0 \\ 3-10 & -9-0 \end{bmatrix}$$

$$BA = \begin{bmatrix} 3 & 3 \\ -7 & -9 \end{bmatrix}$$

Hence proved $AB \neq BA$

(ii) $A(BC) = (AB)C$

Sol. Associative law of Multiplication

$$A(BC) = (AB)C \quad \text{L.H.S.}$$

Firstly

$$BC = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$BC = \begin{bmatrix} (1)(2)+(2)(1) & (1)(1)+(2)(3) \\ (-3)(2)+(-5)(1) & (-3)(1)+(-5)(3) \end{bmatrix}$$

$$BC = \begin{bmatrix} 2+2 & 1+6 \\ -6-5 & -3-15 \end{bmatrix}$$

$$BC = \begin{bmatrix} 4 & 7 \\ -11 & -18 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 4 & 7 \\ -11 & -18 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} (-1)(4)+(3)(-11) & (-1)(7)+(3)(-18) \\ (2)(4)+(0)(-11) & (2)(7)+(0)(-18) \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} -4-33 & -7-54 \\ 8-0 & 14-0 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} -37 & -61 \\ 8 & 14 \end{bmatrix}$$

R.H.S.

$$AB = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$$

$$AB = \begin{bmatrix} (-1)(1)+(3)(-3) & (-1)(2)+(3)(-5) \\ (2)(1)+(0)(-3) & (2)(2)+(0)(-5) \end{bmatrix}$$

$$AB = \begin{bmatrix} -1-9 & -2-15 \\ 2-0 & 4-0 \end{bmatrix} = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}$$

$$(AB)C = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$(AB)C = \begin{bmatrix} (-10)(2)+(-17)(1) & (-10)(1)+(-17)(3) \\ (2)(2)+(4)(1) & (2)(1)+(4)(3) \end{bmatrix}$$

$$(AB)C = \begin{bmatrix} -20-17 & -10-51 \\ 4+4 & 2+12 \end{bmatrix}$$

$$(AB)C = \begin{bmatrix} -37 & -61 \\ 8 & 14 \end{bmatrix}$$

Hence proved $L.H.S = R.H.S$

(iii) $A(B + C) = AB + AC$

Sol. Distributive Law of Multiplication over addition.

L.H.S.

$A(B + C)$

$$B + C = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$B + C = \begin{bmatrix} 1+2 & 2+1 \\ -3+1 & -5+3 \end{bmatrix}$$

$$B + C = \begin{bmatrix} 3 & 3 \\ -2 & -2 \end{bmatrix}$$

$$A(B + C) = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 3 & 3 \\ -2 & -2 \end{bmatrix}$$

$$A(B + C) = \begin{bmatrix} (-1)(3)+(3)(-2) & (-1)(3)+(3)(-2) \\ (2)(3)+(0)(-2) & (2)(3)+(0)(-2) \end{bmatrix}$$

$$A(B + C) = \begin{bmatrix} -3-6 & -3-6 \\ 6-0 & 6-0 \end{bmatrix}$$

$$A(B + C) = \begin{bmatrix} -9 & -9 \\ 6 & 6 \end{bmatrix}$$

R.H.S

$$AB = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$$

$$AB = \begin{bmatrix} (-1)(1)+(3)(-3) & (-1)(2)+(3)(-5) \\ (2)(1)+(0)(-3) & (2)(2)+(0)(-5) \end{bmatrix}$$

$$AB = \begin{bmatrix} -1-9 & -2-15 \\ 2-0 & 4-0 \end{bmatrix} = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}$$

$$AC = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$AC = \begin{bmatrix} (-1)(2)+(3)(1) & (-1)(1)+(3)(3) \\ (2)(2)+(0)(1) & (2)(1)+(0)(3) \end{bmatrix}$$

$$AC = \begin{bmatrix} -2+3 & -1+9 \\ 4+0 & 2+0 \end{bmatrix}$$

$$AC = \begin{bmatrix} 1 & 8 \\ 2 & 4 \end{bmatrix}$$

$$AB + AC = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 8 \\ 2 & 4 \end{bmatrix}$$

$$AB + AC = \begin{bmatrix} -10+1 & -17+8 \\ 2+4 & 4+2 \end{bmatrix}$$

$$AB + AC = \begin{bmatrix} -9 & -9 \\ 6 & 6 \end{bmatrix}$$

Hence proved L.H.S = R.H.S**(iv) $A(B - C) = AB - AC$** **Sol. Distributive Law of Multiplication over Subtraction.****L.H.S.** **$A(B - C)$**

$$B - C = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} - \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$B - C = \begin{bmatrix} 1-2 & 2-1 \\ -3-1 & -5-3 \end{bmatrix}$$

$$B - C = \begin{bmatrix} -1 & 1 \\ -4 & -8 \end{bmatrix}$$

$$A(B - C) = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ -4 & -8 \end{bmatrix}$$

$$A(B - C) = \begin{bmatrix} (-1)(-1)+(3)(-4) & (-1)(1)+(3)(-8) \\ (2)(-1)+(0)(-4) & (2)(1)+(0)(-8) \end{bmatrix}$$

$$A(B - C) = \begin{bmatrix} 1-12 & -1-24 \\ -2-0 & 2-0 \end{bmatrix}$$

$$A(B - C) = \begin{bmatrix} -11 & -25 \\ -2 & 2 \end{bmatrix}$$

R.H.S: **$AB - AC$**

$$AB = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$$

$$AB = \begin{bmatrix} (-1)(1)+(3)(-3) & (-1)(2)+(3)(-5) \\ (2)(1)+(0)(-3) & (2)(2)+(0)(-5) \end{bmatrix}$$

$$AB = \begin{bmatrix} -1-9 & -2-15 \\ 2-0 & 4-0 \end{bmatrix}$$

$$AB = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}$$

$$AC = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$AC = \begin{bmatrix} (-1)(2)+(3)(1) & (-1)(1)+(3)(3) \\ (2)(2)+(0)(1) & (2)(1)+(0)(3) \end{bmatrix}$$

$$AC = \begin{bmatrix} -2+3 & -1+9 \\ 4+0 & 2+0 \end{bmatrix}$$

$$AC = \begin{bmatrix} 1 & 8 \\ 4 & 2 \end{bmatrix}$$

$$AB - AC = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix} - \begin{bmatrix} 1 & 8 \\ 4 & 2 \end{bmatrix}$$

$$AB - AC = \begin{bmatrix} -10-1 & -17-8 \\ 2-4 & 4-2 \end{bmatrix}$$

$$AB - AC = \begin{bmatrix} -11 & -25 \\ -2 & 2 \end{bmatrix}$$

Hence proved L.H.S = R.H.S**Q.6: For the matrices $A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$,**

$$B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}, C = \begin{bmatrix} -2 & 6 \\ 3 & -9 \end{bmatrix}$$

verify that:

$$(i) (AB)^t = B^t A^t \quad (ii) (BC)^t = C^t B^t$$

$$(i) (AB)^t = B^t A^t$$

L.H.S. $(AB)^t$

$$\text{Sol. } A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$$

$$AB = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$$

$$AB = \begin{bmatrix} (-1)(1)+(3)(-3) & (-1)(2)+(3)(-5) \\ (2)(1)+(0)(-3) & (2)(2)+(0)(-5) \end{bmatrix}$$

$$AB = \begin{bmatrix} -1-9 & -2-15 \\ 2-0 & 4-0 \end{bmatrix}$$

$$AB = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}$$

$$(AB)^t = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}^t$$

$$(AB)^t = \begin{bmatrix} -10 & 2 \\ -17 & 4 \end{bmatrix}$$

$$\text{R.H.S} = B^t A^t$$

$$B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}, A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix},$$

$$B^t = \begin{bmatrix} 1 & -3 \\ 2 & -5 \end{bmatrix}, A^t = \begin{bmatrix} -1 & 2 \\ 3 & 0 \end{bmatrix}$$

$$B^t A^t = \begin{bmatrix} 1 & -3 \\ 2 & -5 \end{bmatrix} \begin{bmatrix} -1 & 2 \\ 3 & 0 \end{bmatrix}$$

$$B^t A^t = \begin{bmatrix} (1)(-1)+(-3)(3) & (1)(2)+(-3)(0) \\ (2)(-1)+(-5)(3) & (2)(2)+(-5)(0) \end{bmatrix}$$

$$B^t A^t = \begin{bmatrix} -1-9 & 2-0 \\ -2-15 & 4-0 \end{bmatrix}$$

$$B^t A^t = \begin{bmatrix} -10 & 2 \\ -17 & 4 \end{bmatrix}$$

Hence proved L.H.S = R.H.S

(ii) $(BC)^t = C^t B^t$

Sol. $(BC)^t = C^t B^t$

L.H.S

$$BC = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} \begin{bmatrix} -2 & 6 \\ 3 & -9 \end{bmatrix}$$

$$BC = \begin{bmatrix} (1)(-2)+(2)(3) & (1)(6)+(2)(-9) \\ (-3)(-2)+(-5)(3) & (-3)(6)+(-5)(-9) \end{bmatrix}$$

$$BC = \begin{bmatrix} -2+6 & 6-18 \\ 6-15 & -18+45 \end{bmatrix}$$

$$BC = \begin{bmatrix} 4 & -12 \\ -9 & 27 \end{bmatrix}$$

$$(BC)^t = \begin{bmatrix} 4 & -12 \\ -9 & 27 \end{bmatrix}^t$$

$$(BC)^t = \begin{bmatrix} 4 & -9 \\ -12 & 27 \end{bmatrix}$$

R.H.S

$$C^t B^t$$

$$C^t = \begin{bmatrix} -2 & 3 \\ 6 & -9 \end{bmatrix}, B^t = \begin{bmatrix} 1 & -3 \\ 2 & -5 \end{bmatrix}$$

$$C^t B^t = \begin{bmatrix} -2 & 3 \\ 6 & -9 \end{bmatrix} \begin{bmatrix} 1 & -3 \\ 2 & -5 \end{bmatrix}$$

$$C^t B^t = \begin{bmatrix} (-2)(1)+(3)(2) & (-2)(-3)+(3)(-5) \\ (6)(1)+(-9)(2) & (6)(-3)+(-9)(-5) \end{bmatrix}$$

$$C^t B^t = \begin{bmatrix} -2+6 & 6-15 \\ 6-18 & -18+45 \end{bmatrix}$$

$$C^t B^t = \begin{bmatrix} 4 & -9 \\ -12 & 27 \end{bmatrix}$$

Hence proved L.H.S = R.H.S

Determinant:

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be a 2-by-2 square matrix.

The determinant of A is denoted by $\det A$ or $|A|$ is defined as

$$|A| = \det A = \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - cb = \lambda \in \mathbb{R}$$

Example:

$$B = \begin{bmatrix} 1 & 3 \\ 2 & -2 \end{bmatrix}$$

Sol. $B = \begin{bmatrix} 1 & 3 \\ 2 & -2 \end{bmatrix}$

$$|B| = \begin{vmatrix} 1 & 3 \\ 2 & -2 \end{vmatrix} = (1)(-2) - (2)(3) \\ = -2 - 6 = -8$$

Singular matrix:

A square matrix is singular if its determinant is zero. e.g; see $|A| = 0$

$$|A| = \begin{vmatrix} 3 & 6 \\ 2 & 4 \end{vmatrix} = (3)(4) - (2)(6) \\ = 12 - 12 = 0$$

Non-singular matrix:

A square matrix is non-singular if its determinant is not equal to zero. e.g; See $|C| \neq 0$

$$|C| = \begin{vmatrix} 7 & -9 \\ 3 & 5 \end{vmatrix} = (7)(5) - (3)(-9)$$

$= 35 + 27 = 62 \neq 0$ C is a non-singular matrix.

Adjoint of a matrix:

Adjoint of a square matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is

obtained by interchanging the diagonal entries and changing the signs of other entries Ad joint of matrix A is denoted as Adj A

$$\text{Adj } A = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

EXERCISE 1.5

Q.1: Find the determinant of the following matrices.

Note: A determinant is denoted by $\det A$ or $|A|$

(i) $A = \begin{bmatrix} -1 & 1 \\ 2 & 0 \end{bmatrix}$

Sol. $A = \begin{bmatrix} -1 & 1 \\ 2 & 0 \end{bmatrix}$

$$|A| = \begin{vmatrix} -1 & 1 \\ 2 & 0 \end{vmatrix}$$

$$|A| = (-1)(0) - (2)(1)$$

$$|A| = 0 - 2 = -2$$

(ii) $B = \begin{bmatrix} 1 & 3 \\ 2 & -2 \end{bmatrix}$

Sol. $B = \begin{bmatrix} 1 & 3 \\ 2 & -2 \end{bmatrix}$

$$|B| = \begin{vmatrix} 1 & 3 \\ 2 & -2 \end{vmatrix}$$

$$|B| = (1)(-2) - (2)(3)$$

$$|B| = -2 - 6 = -8$$

(iii) $C = \begin{bmatrix} 3 & 2 \\ 3 & 2 \end{bmatrix}$

Sol. $C = \begin{bmatrix} 3 & 2 \\ 3 & 2 \end{bmatrix}$

$$|C| = \begin{vmatrix} 3 & 2 \\ 3 & 2 \end{vmatrix}$$

$$|C| = (3)(2) - (3)(2)$$

$$|C| = 6 - 6 = 0$$

(iv) $D = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$

Sol. $D = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$

$$|D| = (3)(4) - (1)(2)$$

$$|D| = 12 - 2 = 10$$

Q.2: Find which of the following matrices are singular or non-singular?

(i) $A = \begin{bmatrix} 3 & 6 \\ 2 & 4 \end{bmatrix}$

Sol. $A = \begin{bmatrix} 3 & 6 \\ 2 & 4 \end{bmatrix}$

$$|A| = \begin{vmatrix} 3 & 6 \\ 2 & 4 \end{vmatrix}$$

$$|A| = (3)(4) - (2)(6)$$

$$|A| = 12 - 12 = 0$$

A is a singular matrix because its determinant = 0

(ii) $B = \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix}$

Sol. $B = \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix}$

$$|B| = \begin{vmatrix} 4 & 1 \\ 3 & 2 \end{vmatrix}$$

$$|B| = (4)(2) - (3)(1)$$

$$|B| = 8 - 3 \neq 0$$

B is a non singular matrix because its determinant is $\neq 0$

(iii) $C = \begin{bmatrix} 7 & -9 \\ 3 & 5 \end{bmatrix}$

Sol. $C = \begin{bmatrix} 7 & -9 \\ 3 & 5 \end{bmatrix}$

$$|C| = \begin{vmatrix} 7 & -9 \\ 3 & 5 \end{vmatrix}$$

$$|C| = (7)(5) - (3)(-9)$$

$$|C| = 35 + 27 = 62$$

C is a non-singular matrix because its determinant is $\neq 0$

(iv) $D = \begin{bmatrix} 5 & -10 \\ -2 & 4 \end{bmatrix}$

Sol. $D = \begin{bmatrix} 5 & -10 \\ -2 & 4 \end{bmatrix}$

$$|D| = \begin{vmatrix} 5 & -10 \\ -2 & 4 \end{vmatrix}$$

$$|D| = (5)(4) - (-2)(-10)$$

$$|D| = 20 - 20 = 0$$

D is a singular matrix because its determinant = 0

Q.3: Find the multiplicative inverse (if it exists) of each.

(i) $A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$

Sol. $A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$

$$|A| = \begin{vmatrix} -1 & 3 \\ 2 & 0 \end{vmatrix}$$

$$|A| = (-1)(0) - (2)(3)$$

$$|A| = 0 - 6 = -6 \neq 0$$

A is a Non-singular matrix so solution possible.

$$\text{Adj } A = \begin{bmatrix} 0 & -3 \\ -2 & -1 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 0 & -3 \\ -2 & -1 \end{bmatrix}}{-6}$$

$$A^{-1} = -\frac{1}{6} \begin{bmatrix} 0 & -3 \\ -2 & -1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} (-\frac{1}{6})(0) & (-\frac{1}{6})(-3) \\ (-\frac{1}{6})(-2) & (-\frac{1}{6})(-1) \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 0 & \frac{1}{2} \\ \frac{1}{3} & \frac{1}{6} \end{bmatrix}$$

(ii) $B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$

Sol. $B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$

$$|B| = \begin{vmatrix} 1 & 2 \\ -3 & -5 \end{vmatrix}$$

$$|B| = (1)(-5) - (-3)(2)$$

$$|B| = -5 + 6 \\ = 1 \neq 0$$

B is a Non singular matrix so solution possible.

$$\text{Adj } B = \begin{bmatrix} -5 & -2 \\ 3 & 1 \end{bmatrix}$$

$$B^{-1} = \frac{\text{Adj } B}{|B|}$$

$$B^{-1} = \frac{\begin{bmatrix} -5 & -2 \\ 3 & 1 \end{bmatrix}}{1}$$

$$B^{-1} = \begin{bmatrix} -5 & -2 \\ 3 & 1 \end{bmatrix}$$

(iii) $C = \begin{bmatrix} -2 & 6 \\ 3 & -9 \end{bmatrix}$

Sol. $C = \begin{bmatrix} -2 & 6 \\ 3 & -9 \end{bmatrix}$

$$|C| = \begin{vmatrix} -2 & 6 \\ 3 & -9 \end{vmatrix}$$

$$|C| = (-2)(-9) - (3)(6)$$

$$|C| = 18 - 18 = 0$$

C is singular matrix so multiplicative inverse does not exist.

(iv) $D = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 1 & 2 \end{bmatrix}$

Sol. $D = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 1 & 2 \end{bmatrix}$

$$|D| = \begin{vmatrix} 1 & 3 \\ 2 & 4 \\ 1 & 2 \end{vmatrix}$$

$$D = \left(\frac{1}{2} \right) (2) - (1) \left(\frac{3}{4} \right)$$

$$D = 1 - \frac{3}{4}$$

$$D = \frac{4-3}{4} = \frac{1}{4} \neq 0$$

D is a Non singular matrix so solution possible.

$$D = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 1 & 2 \end{bmatrix}$$

$$\text{Adj } D = \begin{bmatrix} 2 & -3 \\ -1 & \frac{1}{2} \end{bmatrix}$$

$$D^{-1} = \frac{\text{Adj } D}{|D|}$$

$$D^{-1} = \frac{\begin{bmatrix} 2 & -3 \\ -1 & \frac{1}{2} \end{bmatrix}}{\frac{1}{4}}$$

$$D^{-1} = \frac{1}{4}$$

$$D^{-1} = 4 \begin{bmatrix} 2 & -3 \\ -1 & \frac{1}{2} \end{bmatrix}$$

$$D^{-1} = \begin{bmatrix} (4)(2) & (4)\left(\frac{-3}{4}\right) \\ (-1)(4) & \left(\frac{1}{2}\right)(4) \end{bmatrix}$$

$$D^{-1} = \begin{bmatrix} 8 & -3 \\ -4 & 2 \end{bmatrix}$$

Q.4: If $A = \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & -1 \\ 2 & -2 \end{bmatrix}$ then

(i) $A(\text{Adj } A) = (\text{Adj } A)A = (\det A)I$

(ii) $BB^{-1} = I = B^{-1}B$

(i) $A(\text{Adj } A) = (\text{Adj } A)A = (\det A)I$

Sol. $A = \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix}$

$\text{Adj } A = \begin{bmatrix} 6 & -2 \\ -4 & 1 \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix} \begin{bmatrix} 6 & -2 \\ -4 & 1 \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} (1)(6)+(2)(-4) & (1)(-2)+(2)(1) \\ (4)(6)+(6)(-4) & (4)(-2)+(6)(1) \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} 6-8 & -2+2 \\ 24-24 & -8+6 \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix}$

$(\text{Adj } A)A = A(\text{Adj } A)$

$A(\text{Adj } A) = \begin{bmatrix} 6 & -2 \\ -4 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} (6)(1)+(-2)(4) & (6)(2)+(-2)(6) \\ (-4)(1)+(1)(4) & (-4)(2)+(1)(6) \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} 6-8 & 12-12 \\ -4+4 & -8+6 \end{bmatrix}$

$A(\text{Adj } A) = \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix}$

$(\det A) I \quad \therefore \quad \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$(\det A) = \begin{vmatrix} 1 & 2 \\ 4 & 6 \end{vmatrix}$

$|A| = \begin{vmatrix} 1 & 2 \\ 4 & 6 \end{vmatrix}$

$|A| = (1)(6) - (4)(2)$

$|A| = 6 - 8 = -2$

$(\det A)I = -2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$(\det A)I = \begin{bmatrix} (-2)(1) & (-2)(0) \\ (-2)(0) & (-2)(1) \end{bmatrix}$

$(\det A)I = \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix}$

It is proved $A(\text{Adj } A) = (\text{Adj } A)A = (\det A)I$

(ii) $BB^{-1} = I = B^{-1}B$

Sol.

R.H.S $B^{-1}B = I$

$B = \begin{bmatrix} 3 & -1 \\ 2 & -2 \end{bmatrix}$

$|B| = \begin{vmatrix} 3 & -1 \\ 2 & -2 \end{vmatrix}$

$= (3)(-2) - (2)(-1) = -6 + 2 = -4 \neq 0$

B is a Non-singular matrix so solution is possible.

$\text{Adj } B = \begin{bmatrix} -2 & 1 \\ -2 & 3 \end{bmatrix}$

$B^{-1} = \frac{1}{|B|} \text{Adj } B$

$B^{-1} = \frac{1}{-4} \begin{bmatrix} -2 & 1 \\ -2 & 3 \end{bmatrix}$

$B^{-1}B = \frac{1}{-4} \begin{bmatrix} -2 & 1 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 3 & -1 \\ 2 & -2 \end{bmatrix}$

$B^{-1}B = \frac{1}{-4} \begin{bmatrix} (-2)(3)+(1)(2) & (-2)(-1)+(1)(-2) \\ (-2)(3)+(3)(2) & (-2)(-1)+(3)(-2) \end{bmatrix}$

$B^{-1}B = \frac{1}{-4} \begin{bmatrix} -6+2 & 2-2 \\ -6+6 & 2-6 \end{bmatrix}$

$B^{-1}B = \frac{1}{-4} \begin{bmatrix} -4 & 0 \\ 0 & -4 \end{bmatrix}$

$B^{-1}B = \begin{bmatrix} \frac{-4}{-4} & \frac{0}{-4} \\ \frac{0}{-4} & \frac{-4}{-4} \end{bmatrix}$

$B^{-1}B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$B^{-1}B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \therefore \quad I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

I = B⁻¹B

L.H.S

$BB^{-1} = \frac{1}{-4} \begin{bmatrix} 3 & -1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} -2 & 1 \\ -2 & 3 \end{bmatrix}$

$BB^{-1} = \frac{1}{-4} \begin{bmatrix} (3)(-2)+(-1)(-2) & (3)(1)+(-1)(3) \\ (2)(-2)+(-2)(-2) & (2)(1)+(-2)(3) \end{bmatrix}$

$BB^{-1} = \frac{1}{-4} \begin{bmatrix} -6+2 & 3-3 \\ -4+4 & 2-6 \end{bmatrix}$

$BB^{-1} = \frac{1}{-4} \begin{bmatrix} -4 & 0 \\ 0 & -4 \end{bmatrix}$

$BB^{-1} = \begin{bmatrix} \frac{-4}{-4} & \frac{0}{-4} \\ \frac{0}{-4} & \frac{-4}{-4} \end{bmatrix}$

$BB^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$BB^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Hence proved $BB^{-1} = I = B^{-1}B$

Q.5: Determine whether the given matrices are multiplicative inverses of each other.

(i) $\begin{bmatrix} 3 & 5 \\ 4 & 7 \end{bmatrix}$ and $\begin{bmatrix} 7 & -5 \\ -4 & 3 \end{bmatrix}$

Sol.
$$= \begin{bmatrix} 3 & 5 \\ 4 & 7 \end{bmatrix} \begin{bmatrix} 7 & -5 \\ -4 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} (3)(7)+(5)(-4) & (3)(-5)+(5)(3) \\ (4)(7)+(7)(-4) & (4)(-5)+(7)(3) \end{bmatrix}$$

$$= \begin{bmatrix} 21-20 & -15+15 \\ 28-28 & -20+21 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \mathbf{I}$$

So both matrices are multiplicative inverse of each other.

(ii) $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$ and $\begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix}$

Sol.
$$= \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(-3)+(2)(2) & (1)(2)+(2)(-1) \\ (2)(-3)+(3)(2) & (2)(2)+(3)(-1) \end{bmatrix}$$

$$= \begin{bmatrix} -3+4 & 2-2 \\ -6+6 & 4-3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \Rightarrow \mathbf{I}$$

So both matrices are multiplicative inverse of each other.

Q.6: If $A = \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} -4 & -2 \\ 1 & -1 \end{bmatrix}$,

$D = \begin{bmatrix} 3 & 1 \\ -2 & 2 \end{bmatrix}$ then verify that

(i) $(AB)^{-1} = B^{-1}A^{-1}$

(ii) $(DA)^{-1} = A^{-1}D^{-1}$

(i) $(AB)^{-1} = B^{-1}A^{-1}$

Sol.

L.H.S

$$AB = \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} -4 & -2 \\ 1 & -1 \end{bmatrix} =$$

$$AB = \begin{bmatrix} (4)(-4)+(0)(1) & (4)(-2)+(0)(-1) \\ (-1)(-4)+(2)(1) & (-1)(-2)+(2)(-1) \end{bmatrix}$$

$$AB = \begin{bmatrix} -16+0 & -8+0 \\ 4+2 & 2-2 \end{bmatrix} = \begin{bmatrix} -16 & -8 \\ 6 & 0 \end{bmatrix}$$

$$|AB| = \begin{vmatrix} -16 & -8 \\ 6 & 0 \end{vmatrix}$$

$$AB = (-16)(0) - (6)(-8) = 0 + 48$$

$$|AB| = 48$$

$$\text{Adj } AB = \begin{bmatrix} 0 & 8 \\ -6 & -16 \end{bmatrix}$$

$$(AB)^{-1} = \frac{\text{Adj } AB}{|AB|}$$

$$(AB)^{-1} = \frac{\begin{bmatrix} 0 & 8 \\ -6 & -16 \end{bmatrix}}{48}$$

$$(AB)^{-1} = \frac{1}{48} \begin{bmatrix} 0 & 8 \\ -6 & -16 \end{bmatrix}$$

$$(AB)^{-1} = \begin{bmatrix} (\frac{1}{48})(0) & (\frac{1}{48})(8) \\ (\frac{1}{48})(-6) & (\frac{1}{48})(-16) \end{bmatrix}$$

$$(AB)^{-1} = \begin{bmatrix} 0 & \frac{1}{6} \\ -\frac{1}{8} & -\frac{1}{3} \end{bmatrix}$$

R.H.S

$$A = \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 4 & 0 \\ -1 & 2 \end{vmatrix}$$

$$|A| = (4)(2) - (-1)(0)$$

$$|A| = 8 - 0$$

$$|A| = 8$$

$$\text{Adj } A = \begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$A^{-1} = \frac{\begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}}{8} = \frac{1}{8} \begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} (2)(\frac{1}{8}) & (0)(\frac{1}{8}) \\ (1)(\frac{1}{8}) & (4)(\frac{1}{8}) \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{1}{2} \end{bmatrix}$$

$$B = \begin{bmatrix} -4 & -2 \\ 1 & -1 \end{bmatrix}$$

$$|B| = \begin{vmatrix} -4 & -2 \\ 1 & -1 \end{vmatrix}$$

$$= (-4)(-1) - (1)(-2)$$

$$|B| = 4 + 2 = 6$$

$$\text{Adj } B = \begin{bmatrix} -1 & 2 \\ -1 & -4 \end{bmatrix}$$

$$B^{-1} = \frac{\text{Adj } B}{|B|}$$

$$B^{-1} = \frac{\begin{bmatrix} -1 & 2 \\ -1 & -4 \end{bmatrix}}{6}$$

$$B^{-1} = \frac{1}{6} \begin{bmatrix} -1 & 2 \\ -1 & -4 \end{bmatrix}$$

$$B^{-1} = \begin{bmatrix} (\frac{1}{6})(-1) & (2)(\frac{1}{6}) \\ (\frac{1}{6})(-1) & (-4)(\frac{1}{6}) \end{bmatrix}$$

$$B^{-1} = \begin{bmatrix} -\frac{1}{6} & \frac{1}{3} \\ -\frac{1}{6} & -\frac{2}{3} \end{bmatrix}$$

$$B^{-1}A^{-1} = \begin{bmatrix} -\frac{1}{6} & \frac{1}{3} \\ -\frac{1}{6} & -\frac{2}{3} \end{bmatrix} \begin{bmatrix} \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{1}{2} \end{bmatrix}$$

$$B^{-1}A^{-1} = \begin{bmatrix} (-\frac{1}{6})(\frac{1}{4})+(\frac{1}{3})(\frac{1}{8}) & (-\frac{1}{6})(0)+(\frac{1}{3})(\frac{1}{2}) \\ (-\frac{1}{6})(\frac{1}{4})+(-\frac{2}{3})(\frac{1}{8}) & (-\frac{1}{6})(0)+(-\frac{2}{3})(\frac{1}{2}) \end{bmatrix}$$

$$B^{-1}A^{-1} = \begin{bmatrix} -\frac{1}{24}+\frac{1}{24} & 0+\frac{1}{6} \\ -\frac{1}{24}-\frac{2}{24} & 0-\frac{2}{6} \end{bmatrix}$$

$$B^{-1}A^{-1} = \begin{bmatrix} 0 & \frac{1}{6} \\ -\frac{3}{24} & -\frac{1}{3} \end{bmatrix} \Rightarrow B^{-1}A^{-1} = \begin{bmatrix} 0 & \frac{1}{6} \\ -\frac{1}{8} & -\frac{1}{3} \end{bmatrix}$$

Hence proved

L.H.S = R.H.S

(ii) $(DA)^{-1} = A^{-1}D^{-1}$

$$D = \begin{bmatrix} 3 & 1 \\ -2 & 2 \end{bmatrix}$$

$$A = \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix}$$

Sol.

L.H.S

$$(DA) = \begin{bmatrix} 3 & 1 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix}$$

$$(DA) = \begin{bmatrix} (3)(4)+(1)(-1) & (3)(0)+(1)(2) \\ (-2)(4)+(2)(-1) & (-2)(0)+(2)(2) \end{bmatrix}$$

$$(DA) = \begin{bmatrix} 12-1 & 0+2 \\ -8-2 & 0+4 \end{bmatrix}$$

$$(DA) = \begin{bmatrix} 11 & 2 \\ -10 & 4 \end{bmatrix}$$

$$|DA| = \begin{vmatrix} 11 & 2 \\ -10 & 4 \end{vmatrix}$$

$$|DA| = (11)(4) - (-10)(2) = 44 + 20$$

$$|DA| = 64$$

$$\text{Adj } DA = \begin{bmatrix} 4 & -2 \\ 10 & 11 \end{bmatrix}$$

$$(DA)^{-1} = \frac{(\text{Adj } DA)}{|DA|}$$

$$(DA)^{-1} = \frac{\begin{bmatrix} 4 & -2 \\ 10 & 11 \end{bmatrix}}{64}$$

$$(DA)^{-1} = \frac{1}{64} \begin{bmatrix} 4 & -2 \\ 10 & 11 \end{bmatrix}$$

$$(DA)^{-1} = \begin{bmatrix} (\frac{1}{64})(4) & (\frac{1}{64})(-2) \\ (\frac{1}{64})(10) & (\frac{1}{64})(11) \end{bmatrix}$$

$$(DA)^{-1} = \begin{bmatrix} \frac{1}{16} & -\frac{1}{32} \\ \frac{5}{32} & \frac{11}{64} \end{bmatrix}$$

R.H.S:

$A^{-1}D^{-1}$

$$A = \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 4 & 0 \\ -1 & 2 \end{vmatrix}$$

$$|A| = (4)(2) - (-1)(0)$$

$$|A| = 8 - 0 = 8$$

$$\text{Adj } A = \begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$A^{-1} = \frac{\begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}}{8}$$

$$A^{-1} = \frac{1}{8} \begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} (\frac{1}{8})(2) & (\frac{1}{8})(0) \\ (\frac{1}{8})(1) & (\frac{1}{8})(4) \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{1}{2} \end{bmatrix}$$

$$|D| = \begin{vmatrix} 3 & 1 \\ -2 & 2 \end{vmatrix}$$

$$|D| = (3)(2) - (-2)(1)$$

$$|D| = 6 + 2 = 8$$

$$\text{Adj}D = \begin{bmatrix} 2 & -1 \\ 2 & 3 \end{bmatrix}$$

$$D^{-1} = \frac{\text{Adj}D}{|D|}$$

$$D^{-1} = \frac{\begin{bmatrix} 2 & -1 \\ 2 & 3 \end{bmatrix}}{8}$$

$$D^{-1} = \begin{bmatrix} (\frac{1}{8})(2) & (\frac{1}{8})(-1) \\ (\frac{1}{8})(2) & (\frac{1}{8})(3) \end{bmatrix}$$

$$D^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{8} \\ \frac{1}{4} & \frac{3}{8} \end{bmatrix}$$

$$A^{-1}D^{-1} = \begin{bmatrix} \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} \frac{1}{4} & -\frac{1}{8} \\ \frac{1}{4} & \frac{3}{8} \end{bmatrix}$$

$$A^{-1}D^{-1} = \begin{bmatrix} (\frac{1}{4})(\frac{1}{4}) + (0)(\frac{1}{4}) & (\frac{1}{4})(-\frac{1}{8}) + (0)(\frac{3}{8}) \\ (\frac{1}{8})(\frac{1}{4}) + (\frac{1}{2})(\frac{1}{4}) & (\frac{1}{8})(-\frac{1}{8}) + (\frac{1}{2})(\frac{3}{8}) \end{bmatrix}$$

$$A^{-1}D^{-1} = \begin{bmatrix} \frac{1}{16} + 0 & \frac{-1}{32} + 0 \\ \frac{1}{32} + \frac{1}{8} & \frac{-1}{64} + \frac{3}{16} \end{bmatrix}$$

$$A^{-1}D^{-1} = \begin{bmatrix} \frac{1}{16} & \frac{-1}{32} \\ \frac{1+4}{32} & \frac{-1+12}{64} \end{bmatrix} \quad A^{-1}D^{-1} = \begin{bmatrix} \frac{1}{16} & \frac{-1}{32} \\ \frac{5}{32} & \frac{11}{64} \end{bmatrix}$$

Hence proved L.H.S = R.H.S

Simultaneous equations:

A system of equation having a common solution is called a system of simultaneous equations

$$ax+by = m$$

$$cx+dy = n$$

where a,b,c,d,m and n are real number

x and y are two variable so these are two variable linear equations

EXERCISE 1.6

Q.1: Use matrices, if possible, to solve the following systems of linear equations by:

(i) The matrix inverse method

(ii) The Cramer's rule.

(i) $2x - 2y = 4$

$3x + 2y = 6$

Sol. Matrix inverse method

By writing in matrix form

$$\begin{bmatrix} 2 & -2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 6 \end{bmatrix} \dots (i)$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 2 & -2 \\ 3 & 2 \end{vmatrix}$$

$$|A| = (2)(2) - (3)(-2)$$

$$|A| = 4 + 6$$

$$|A| = 10 \neq 0$$

Non-singular matrix so solution is possible

$$\text{Adj} A = \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix}$$

$$X = A^{-1}B$$

$$X = \frac{1}{|A|} \text{Adj} A \times B$$

$$X = \frac{1}{10} \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} (2)(4) + (2)(6) \\ (-3)(4) + (2)(6) \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} 8 + 12 \\ -12 + 12 \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} 20 \\ 0 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{20}{10} \\ \frac{0}{10} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

$$x = 2, y = 0$$

(i) **By Cramer's rule**

$$\begin{bmatrix} 2 & -2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 2 & -2 \\ 3 & 2 \end{vmatrix}$$

$$= (2)(2) - (3)(-2)$$

$$= 4 + 6$$

$$= 10 \neq 0$$

Non-singular matrix so solution is possible

$$A_x = \begin{bmatrix} 4 & -2 \\ 6 & 2 \end{bmatrix}$$

$$x = \frac{|A_x|}{|A|}$$

$$x = \frac{\begin{vmatrix} 4 & -2 \\ 6 & 2 \end{vmatrix}}{10}$$

$$x = \frac{(4)(2) - (6)(-2)}{10}$$

$$x = \frac{8 + 12}{10}$$

$$x = \frac{20}{10} = 2$$

$$A_y = \begin{bmatrix} 2 & 4 \\ 3 & 6 \end{bmatrix}$$

$$y = \frac{|A_y|}{|A|}$$

$$y = \frac{\begin{vmatrix} 2 & 4 \\ 3 & 6 \end{vmatrix}}{10}$$

$$y = \frac{(2)(6) - (3)(4)}{10}$$

$$y = \frac{12 - 12}{10}$$

$$y = \frac{0}{10} = 0$$

x = 2, y = 0

(ii) 2x + y = 3

6x + 5y = 1

Sol. By matrix inverse method:

Writing in the matrix form

$$\begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \dots (i)$$

A X = B

$$|A| = \begin{vmatrix} 2 & 1 \\ 6 & 5 \end{vmatrix}$$

$$|A| = (2)(5) - (6)(1)$$

$$|A| = 10 - 6$$

$$|A| = 4 \neq 0$$

Non-singular matrix so solution is possible

$$\text{Adj } A = \begin{bmatrix} 5 & -1 \\ -6 & 2 \end{bmatrix}$$

$$X = A^{-1}B$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$X = \frac{1}{4} \begin{bmatrix} 5 & -1 \\ -6 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$X = \frac{1}{4} \begin{bmatrix} (5)(3) + (-1)(1) \\ (-6)(3) + (2)(1) \end{bmatrix}$$

$$X = \frac{1}{4} \begin{bmatrix} 15 - 1 \\ -18 + 2 \end{bmatrix}$$

$$X = \frac{1}{4} \begin{bmatrix} 14 \\ -16 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{14}{4} \\ \frac{-16}{4} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{7}{2} \\ -4 \end{bmatrix}$$

x = 7/2, y = -4

By Cramer's rule

$$\begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \dots (i)$$

A X = B

$$|A| = \begin{vmatrix} 2 & 1 \\ 6 & 5 \end{vmatrix}$$

$$|A| = (2)(5) - (6)(1)$$

$$|A| = 10 - 6$$

$$|A| = 4 \neq 0$$

Non-singular matrix so solution is possible

$$A_x = \begin{bmatrix} 3 & 1 \\ 1 & 5 \end{bmatrix}$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 3 & 1 \\ 1 & 5 \end{vmatrix}}{|A|} = \frac{(3)(5) - (1)(1)}{4}$$

$$x = \frac{(3)(5) - (1)(1)}{4}$$

$$x = \frac{15 - 1}{4} = \frac{14}{4}$$

$$x = \frac{7}{2}$$

$$A_y = \begin{bmatrix} 2 & 3 \\ 6 & 1 \end{bmatrix}$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 2 & 3 \\ 6 & 1 \end{vmatrix}}{|A|}$$

$$y = \frac{(2)(1) - (3)(6)}{4} = \frac{2 - 18}{4} = \frac{-16}{4} = -4$$

$$y \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7 \\ 2 \\ -4 \end{bmatrix}$$

$$x = \frac{7}{2}, y = -4$$

$$(iii) 4x + 2y = 8$$

$$3x - y = -1$$

Sol. By inverse method:

Writing in the matrix form

$$\begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ -1 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 4 & 2 \\ 3 & -1 \end{vmatrix}$$

$$|A| = (4)(-1) - (3)(2)$$

$$|A| = -4 - 6$$

$$|A| = -10 \neq 0$$

Non-singular matrix so solution is possible

$$X = A^{-1} B$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$\text{Adj } A = \begin{bmatrix} -1 & -2 \\ -3 & 4 \end{bmatrix}$$

$$X = \frac{1}{-10} \begin{bmatrix} -1 & -2 \\ -3 & 4 \end{bmatrix} \begin{bmatrix} 8 \\ -1 \end{bmatrix}$$

$$X = \frac{1}{-10} \begin{bmatrix} (-1)(8) + (-2)(-1) \\ (-3)(8) + (4)(-1) \end{bmatrix}$$

$$X = \frac{1}{-10} \begin{bmatrix} -8 + 2 \\ -24 - 4 \end{bmatrix}$$

$$X = \frac{1}{-10}$$

$$X = \frac{1}{-10} \begin{bmatrix} -6 \\ -28 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{-6}{-10} \\ \frac{-28}{-10} \end{bmatrix} \quad X = \begin{bmatrix} \frac{3}{5} \\ \frac{14}{5} \end{bmatrix}$$

$$x = \frac{3}{5}, y = \frac{14}{5}$$

Cramer's rule

$$\begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ -1 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 4 & 2 \\ 3 & -1 \end{vmatrix}$$

$$|A| = (4)(-1) - (3)(2)$$

$$|A| = -4 - 6$$

$$|A| = -10 \neq 0$$

Non singular matrix so solution is possible

$$x = \frac{|A_x|}{|A|}$$

$$x = \frac{\begin{vmatrix} 8 & 2 \\ -1 & -1 \end{vmatrix}}{-10}$$

$$X = \frac{(8)(-1) - (-1)(2)}{-10}$$

$$X = \frac{-8 + 2}{-10}$$

$$X = \frac{-6}{-10}$$

$$X = \frac{3}{5}$$

$$y = \frac{|A_y|}{|A|}$$

$$Y = \frac{\begin{vmatrix} 4 & 8 \\ 3 & -1 \end{vmatrix}}{-10}$$

$$y = \frac{(4)(-1) - (3)(8)}{-10}$$

$$y = \frac{-4 - 24}{-10} = \frac{-28}{-10}$$

$$y = \frac{14}{5}$$

$$x = \frac{3}{5}, y = \frac{14}{5}$$

$$(iv) 3x - 2y = -6$$

$$5x - 2y = -10$$

Sol. By inverse method

Writing in the matrix form

$$= \begin{bmatrix} 3 & -2 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -6 \\ -10 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 3 & -2 \\ 5 & -2 \end{vmatrix}$$

$$|A| = (3)(-2) - (-2)(5)$$

$$|A| = -6 + 10 = 4 \neq 0$$

$$X = A^{-1}B$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$X = \frac{1}{4} \begin{bmatrix} -2 & 2 \\ -5 & 3 \end{bmatrix} \begin{bmatrix} -6 \\ -10 \end{bmatrix}$$

$$X = \frac{1}{4} \begin{bmatrix} (-2)(-6) + (2)(-10) \\ (-5)(-6) + (3)(-10) \end{bmatrix}$$

$$X = \frac{1}{4} \begin{bmatrix} 12 - 20 \\ 30 - 30 \end{bmatrix}$$

$$X = \frac{1}{4} \begin{bmatrix} -8 \\ 0 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{-8}{4} \\ \frac{0}{4} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -2 \\ 0 \end{bmatrix}$$

$$x = -2, y = 0$$

Cramer's Rule

$$\begin{bmatrix} 3 & -2 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -6 \\ -10 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 3 & -2 \\ 5 & -2 \end{vmatrix}$$

$$|A| = 3 \cdot -2 - 5 \cdot -2 \neq 0$$

$$|A| = -6 + 10 = 4$$

Non singular matrix so solution possible

$$x = \frac{|Ax|}{|A|}$$

$$x = \frac{\begin{vmatrix} -6 & -2 \\ -10 & -2 \end{vmatrix}}{4}$$

$$x = \frac{(-6)(-2) - (-10)(-2)}{4}$$

$$x = \frac{12 - 20}{4}$$

$$x = \frac{-8}{4}$$

$$x = -2$$

$$y = \frac{|Ay|}{|A|}$$

$$y = \frac{\begin{vmatrix} 3 & -6 \\ 5 & -10 \end{vmatrix}}{4}$$

$$y = \frac{(3)(-10) - (5)(-6)}{4}$$

$$= \frac{-30 + 30}{4}$$

$$= \frac{0}{4} = 0$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -2 \\ 0 \end{bmatrix} \quad x = -2, y = 0$$

(v) $3x - 2y = 4$

$$-6x + 4y = 7$$

Sol. By inverse method:

Writing in the matrix form

$$\begin{bmatrix} 3 & -2 \\ -6 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 3 & -2 \\ -6 & 4 \end{vmatrix}$$

$$= (3)(4) - (-6)(-2)$$

$$= 12 - 12 = 0$$

It is singular matrix so solution set is impossible

(vi) $4x + y = 9$

$$-3x - y = -5$$

Writing in the matrix form

Sol. By inverse method

$$\begin{bmatrix} 4 & 1 \\ -3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ -5 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 4 & 1 \\ -3 & -1 \end{vmatrix}$$

$$= (4)(-1) - (-3)(1)$$

$$= -4 + 3 = -1 \neq 0 \text{ Non singular so}$$

possible

$$\text{Adj } A = \begin{bmatrix} -1 & -1 \\ 3 & 4 \end{bmatrix}$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$X = \frac{1}{-1} \begin{bmatrix} -1 & -1 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 9 \\ -5 \end{bmatrix}$$

$$X = \frac{1}{-1} \left[\begin{array}{l} (-1)(9) + (-1)(-5) \\ (3)(9) + (4)(-5) \end{array} \right]$$

$$X = \frac{1}{-1} \begin{bmatrix} -9+5 \\ 27-20 \end{bmatrix}$$

$$X = \frac{1}{-1} \begin{bmatrix} -4 \\ 7 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{-4}{-1} \\ \frac{7}{-1} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$$

$$\mathbf{x = 4, y = -7}$$

Cramer's Rule

$$\begin{bmatrix} 4 & 1 \\ -3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ -5 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 4 & 1 \\ -3 & -1 \end{vmatrix}$$

$$= (4)(-1) - (-3)(1)$$

$$= -4 + 3 = -1$$

$$x = \frac{|Ax|}{|A|}$$

$$= \frac{\begin{vmatrix} 9 & 1 \\ -5 & -1 \end{vmatrix}}{-1}$$

$$= \frac{(9)(-1) - (-5)(1)}{-1}$$

$$= \frac{-9+5}{-1}$$

$$= \frac{-4}{-1}$$

$$x = 4$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 4 & 9 \\ -3 & -5 \end{vmatrix}}{-1}$$

$$= \frac{(4)(-5) - (-3)(9)}{-1}$$

$$= \frac{-20+27}{-1} = \frac{7}{-1}$$

$$= -7$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$$

$$\mathbf{x = 4, y = -7}$$

(vii) $2x - 2y = 4$

$$-5x - 2y = -10$$

Writing in the matrix form

Sol. By inverse matrix

$$\begin{bmatrix} 2 & -2 \\ -5 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -10 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 2 & -2 \\ -5 & -2 \end{vmatrix}$$

$$= (2)(-2) - (-2)(-5)$$

$$= -4 - 10 = -14$$

Non singular matrix so solution is possible

$$\text{Adj } A = \begin{bmatrix} -2 & 2 \\ 5 & 2 \end{bmatrix}$$

$$X = A^{-1}B$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$= \frac{1}{-14} \begin{bmatrix} -2 & 2 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ -10 \end{bmatrix}$$

$$x = \frac{1}{-14} \left[\begin{array}{l} (-2)(4) + (2)(-10) \\ (5)(4) + (2)(-10) \end{array} \right]$$

$$X = \frac{1}{-14} \begin{bmatrix} -8-20 \\ 20-20 \end{bmatrix}$$

$$X = \frac{1}{-14} \begin{bmatrix} -28 \\ 0 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{-28}{-14} \\ \frac{0}{-14} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

$$\mathbf{x = 2, y = 0}$$

Cramer's Rule

$$\begin{bmatrix} 2 & -2 \\ -5 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -10 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 2 & -2 \\ -5 & -2 \end{vmatrix}$$

$$= (2)(-2) - (-2)(-5)$$

$$= -4 - 10 = -14 \neq 0$$

Non singular matrix so solution possible

$$x = \frac{Ax}{A}$$

$$= \frac{\begin{vmatrix} 4 & -2 \\ -10 & -2 \end{vmatrix}}{-14}$$

$$= \frac{(4)(-2) - (-10)(-2)}{-14}$$

$$= \frac{-8 - 20}{-14}$$

$$\frac{-28}{-14} = 2$$

$$y = \frac{|Ay|}{|A|} = \frac{\begin{vmatrix} 2 & 4 \\ -5 & -10 \end{vmatrix}}{-14}$$

$$= \frac{(2)(-10) - (-5)(4)}{-14} \quad y = 0$$

$$= \frac{-20 + 20}{-14}$$

$$x = 2, y = 0$$

(viii) $3x - 4y = 4$

$$x + 2y = 8$$

Sol. By inverse method

Writing in the matrix form

$$\begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$$

$$A X = B$$

$$|A| = \begin{vmatrix} 3 & -4 \\ 1 & 2 \end{vmatrix} = (3)(2) - (-4)(1) = 6 + 4$$

$= 10 \neq 0$ Non singular so possible

$$\text{Adj } A = \begin{bmatrix} 2 & 4 \\ -1 & 3 \end{bmatrix}$$

$$X = A^{-1}B$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$X = \frac{1}{10} \begin{bmatrix} 2 & 4 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 8 \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} (2)(4) + (4)(8) \\ (-1)(4) + (3)(8) \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} 8 + 32 \\ -4 + 24 \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} 40 \\ 20 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{40}{10} \\ \frac{20}{10} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$$

$$\{x = 4, y = 2\}$$

$$\begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$$

Cramer's Rule

$$|A| = \begin{vmatrix} 3 & -4 \\ 1 & 2 \end{vmatrix} = (3)(2) - (1)(-4) = 6 + 4 = 10$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 4 & -4 \\ 8 & 2 \end{vmatrix}}{10}$$

$$= \frac{(4)(2) - (8)(-4)}{10}$$

$$= \frac{8 + 32}{10}$$

$$= \frac{40}{10}$$

$$= 4$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 3 & 4 \\ 1 & 8 \end{vmatrix}}{10}$$

$$= \frac{(3)(8) - (1)(4)}{10} = \frac{20}{10}$$

$$y = 2$$

2: The length of a rectangle is 4 times its width. The perimeter of the rectangle is 150cm. Find the dimensions of the rectangle.

Sol. Suppose lengths of rectangle

$$\text{Width} = x \text{ cm}$$

$$\text{Length} = y \text{ cm}$$

According to given condition

$$4x = y$$

$$4x - y = 0$$

$$2x + 2y = 150$$

Writing the in matrix form

$$\begin{bmatrix} 4 & -1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

$$A \quad X = B$$

$$\text{Where } A = \begin{bmatrix} 4 & -1 \\ 2 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 4 & -1 \\ 2 & 2 \end{vmatrix}$$

$$= (4)(2) - (2)(-1) = 8 + 2 = 10$$

$$\text{Adj } A = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$A^{-1} = \frac{\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}}{10}$$

$$= \begin{bmatrix} \frac{2}{10} & \frac{1}{10} \\ \frac{-2}{10} & \frac{4}{10} \end{bmatrix}$$

$$X = A^{-1}B$$

$$\begin{bmatrix} \frac{2}{10} & \frac{1}{10} \\ \frac{-2}{10} & \frac{4}{10} \end{bmatrix} \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

$$= \begin{bmatrix} \left(\frac{2}{10}\right)(0) + \left(\frac{1}{10}\right)(150) \\ \left(\frac{-2}{10}\right)(0) + \left(\frac{4}{10}\right)(150) \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 15 \\ 0 & 60 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 15 \\ 60 \end{bmatrix}$$

$$x = 15 \quad y = 60$$

Thus Width = 15cm

Length = 60cm

Cramer's Rule

$$A_x = \begin{bmatrix} 0 & -1 \\ 150 & 2 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 4 & 0 \\ 2 & 150 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 4 & -1 \\ 2 & 2 \end{vmatrix}$$

$$= (4)(2) - (2)(-1) = 8 + 2 = 10 \neq 0$$

$$x = \frac{|A_x|}{|A|}$$

$$= \frac{(0)(2) - (150)(-1)}{10}$$

$$= \frac{0 + 150}{10} = \frac{150}{10}$$

$$x = 15$$

$$y = \frac{|A_y|}{|A|}$$

$$= \frac{\begin{vmatrix} 4 & 0 \\ 2 & 150 \end{vmatrix}}{10}$$

$$= \frac{(4)(150) - (2)(0)}{10} = \frac{600}{10}$$

$$= 60$$

$$x = 15$$

$$y = 60$$

3: Two sides of a rectangle differ by 3.5cm. Find the dimensions of the rectangle if its perimeter is 67cm.

Sol. Suppose

$$\text{Length of a rectangle} = x \text{ cm}$$

$$\text{Width of a rectangle} = y \text{ cm}$$

According to the condition

$$x - y = 3.5$$

$$2x + 2y = 67$$

Writing the in matrix form

$$\begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3.5 \\ 67 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 1 & -1 \\ 2 & 2 \end{vmatrix} = (1)(2) - (2)(-1) = 2 + 2 = 4$$

$$\text{Adj } A = \begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$A^{-1} = \frac{\begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix}}{4}$$

$$= \begin{bmatrix} \frac{2}{4} & \frac{1}{4} \\ \frac{-2}{4} & \frac{1}{4} \end{bmatrix}$$

$$X = A^{-1} B$$

$$= \begin{bmatrix} \frac{2}{4} & \frac{1}{4} \\ \frac{-2}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} 3.5 \\ 67 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{4}(3.5) + \frac{1}{4}(67) \\ \frac{-2}{4}(3.5) + \frac{1}{4}(67) \end{bmatrix}$$

$$= \begin{bmatrix} \frac{7}{4} + \frac{67}{4} \\ \frac{-7}{4} + \frac{67}{4} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{74}{4} \\ \frac{60}{4} \end{bmatrix} = \begin{bmatrix} 18.5 \\ 15 \end{bmatrix}$$

$$x = 18.5$$

$$y = 15$$

$$\text{Length} = 18.5\text{cm}$$

$$\text{Width} = 15\text{cm}$$

Cramer's Rule

$$x - y = 3.5$$

$$2x + 2y = 67$$

$$A = \begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix}$$

$$A_x = \begin{bmatrix} 3.5 & -1 \\ 67 & 2 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 1 & 3.5 \\ 2 & 67 \end{bmatrix}$$

$$|A| = (1)(2) - (-1)(2)$$

$$= 2 + 4$$

$$= 4$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 3.5 & -1 \\ 67 & 2 \end{vmatrix}}{4}$$

$$x = \frac{(3.5)(2) - (67)(-1)}{4}$$

$$x = \frac{7 + 67}{4}$$

$$x = \frac{74}{4}$$

$$y = \frac{|A_y|}{A}$$

$$y = \frac{\begin{vmatrix} 1 & 3.5 \\ 2 & 67 \end{vmatrix}}{4}$$

$$\frac{(1)(67) - (-2)(3.5)}{4} = \frac{67-7}{4}$$

$$\frac{60}{4} = 15$$

$$x = 18.5, y = 5$$

4: The third angle of an isosceles triangle is 16° less than the sum of the two equal angles. Find three angles of the triangle.

Sol. Let x , y be the angles of an isosceles triangle.

Then by given condition

$$y = 2x - 160 \text{ or } 2x - 16 = y$$

$$\text{Or } 2x - y = 16^\circ \dots\dots\dots (i)$$

$$\text{And } 2x + y = 180^\circ \dots\dots\dots (ii)$$

Solve: Sum of three angles of Δ is 180 matrix from above equations.

$$\begin{bmatrix} 2 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 16 \\ 180 \end{bmatrix}$$

$$A \quad X = B$$

$$|A| = \begin{vmatrix} 2 & -1 \\ 2 & 1 \end{vmatrix}$$

$$= (2)(1) - (-2)(-1)$$

$$= 2 + 2 = 4 \neq 0$$

$$\text{Adj } A = \begin{bmatrix} 1 & 1 \\ -2 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$= \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -2 & 2 \end{bmatrix}$$

$$X = A^{-1} \times B$$

$$X = \frac{1}{|A|} \text{Adj } A \times B$$

$$= \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} 16 \\ 180 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} (1)(16) + (1)(180) \\ (-2)(16) + (2)(180) \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} 16 + 180 \\ -32 + 360 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} 196 \\ 328 \end{bmatrix}$$

$$X = \begin{bmatrix} \frac{196}{4} \\ \frac{328}{4} \end{bmatrix}$$

$$X = \begin{bmatrix} 49 \\ 82 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 49 \\ 82 \end{bmatrix}$$

$$X = 49$$

$$Y = 82$$

So unknown angles are 49° , 49° and 82°

Cramer's Rule

$$X = \frac{|A_x|}{|A|}$$

$$x = \frac{\begin{vmatrix} 16 & -1 \\ 180 & 1 \end{vmatrix}}{4} = \frac{(16)(1) - (180)(-1)}{4}$$

$$= \frac{16 + 180}{4} = \frac{196}{4} = 49^\circ$$

$$X = 49^\circ$$

$$y = \frac{|A_y|}{|A|}$$

$$y = \frac{\begin{vmatrix} 2 & 16 \\ 2 & 180 \end{vmatrix}}{4} = \frac{(2)(180) - (16)(2)}{4}$$

$$y = \frac{360 - 32}{4} = \frac{328}{4} = 82^\circ$$

So unknown angles are 49° , 49° and 82°

5: One acute angle of a right triangle is 12° more than twice the other acute angle. Find the acute angles of the right triangle.

$$\text{Sol. } \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -12 \\ 90 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} -12 \\ 90 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix} = 2 \times 1 - (-1 \times 1) = 2 + 1 = 3$$

$$X = A^{-1}B$$

$$X = \frac{\text{Adj}A}{A} \times B$$

$$X = \frac{\begin{bmatrix} 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} -12 \\ 90 \end{bmatrix}}{3}$$

$$X = \frac{\begin{bmatrix} -12 & +90 \\ 12 & +180 \end{bmatrix}}{3} = \frac{\begin{bmatrix} 78 \\ 192 \end{bmatrix}}{3}$$

$$= \begin{bmatrix} \frac{78}{3} \\ \frac{192}{3} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 26 \\ 64 \end{bmatrix}$$

$$\Rightarrow x = 26^\circ, y = 64^\circ$$

So one acute angle = 26°

Other acute angle = 64°

Cramer's rule

$$2x - y = -12$$

$$x + y = 90$$

$$|A| = 3, A_x = \begin{bmatrix} -12 & -1 \\ 90 & 1 \end{bmatrix}$$

$$|A_x| = \begin{vmatrix} -12 & -1 \\ 90 & 1 \end{vmatrix}$$

$$|A_x| = -12 + 90 = 78$$

$$x = \frac{|A_x|}{|A|} = \frac{78}{3} = 26^\circ$$

$$\text{Now } A_y = \begin{vmatrix} 2 & -12 \\ 1 & 90 \end{vmatrix} = 2 \times 90 - (-12 \times 1) \\ = 180 + 12 = 192$$

$$\text{So } A_y = \frac{|A_y|}{|A|} = \frac{192}{3}$$

$$y = 64^\circ$$

so one acute angle = 26°

other acute angle = 64°

6: Two cars that are 600 km apart are moving towards each other. Their speeds differ by 6 km per hour and the cars are 123 km apart after $4\frac{1}{2}$ hours. Find the speed of each car.

Sol. Let the 1st speed of car = x km/hr

2nd speed of car = y km/hr

Difference between speeds = $x - y = 6 \dots \dots (i)$

$$\text{Time} = 4\frac{1}{2} \text{ hours} = \frac{9}{2} + 123 + (y) \times \left(\frac{9}{2}\right)$$

$$\frac{9}{2} + \frac{9y}{2} = 600 - 123 = 477$$

$$\Rightarrow 9x + 9y = 954$$

$$9(x + y) = 954 \Rightarrow (x + y) = \frac{954}{9} = 106$$

$$x + y = 106 \dots \dots (ii)$$

The matrix inverse method $x - y = 6$, $x + y = 106$

Sol.

By writing in matrix form of given equation

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 6 \\ 106 \end{bmatrix}$$

$$A \quad X = B$$

$$\Rightarrow X = A^{-1} B$$

$$\text{And } A^{-1} = \frac{1}{|A|} \text{Adj}A$$

$$|A| = \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix} = (1)(1) - (1)(-1)$$

$$|A| = 1 + 1 = 2$$

As $|A| \neq 0$ so solution is possible.

$$\text{Adj } A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$$

$$= A^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$$

$$X = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 6 \\ 106 \end{bmatrix}$$

$$X = \frac{1}{2} \begin{bmatrix} (1) \times (6) + (1) \times (106) \\ (-1) \times (6) + (1) \times (106) \end{bmatrix}$$

$$X = \frac{1}{2} \begin{bmatrix} 6 + 106 \\ -6 + 106 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 112 \\ 100 \end{bmatrix} = \begin{bmatrix} \frac{112}{2} \\ \frac{100}{2} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 56 \\ 50 \end{bmatrix}$$

Hence $x = 56$, $y = 50$

Cramer's rule

Sol. By writing in matrix form

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}, x = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 6 \\ 106 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix}$$

$$= (1)(1) - (-1)(1)$$

$$= 1 + 1 = 2$$

As $|A| \neq 0$ so solution is possible.

$$\text{Now } |A_x| = \begin{vmatrix} 6 & -1 \\ 106 & 1 \end{vmatrix}$$

$$\Rightarrow |A_x| = \begin{vmatrix} 6 & -1 \\ 106 & 1 \end{vmatrix}$$

$$= (6)(1) - (-1)(106)$$

$$= 6 + 106 = 112$$

$$A_y = \begin{vmatrix} 1 & 6 \\ 1 & 106 \end{vmatrix}$$

$$\Rightarrow |A_y| = \begin{vmatrix} 1 & 6 \\ 1 & 106 \end{vmatrix}$$

$$= (1)(106) - (1)(6)$$

$$= 106 - 6 = 100$$

$$x = \frac{|A_x|}{|A|} = \frac{112}{2} = 56 \text{ and}$$

$$y = \frac{|A_y|}{|A|} = \frac{100}{2} = 50$$

So, $x = 56$, $y = 100$

REVIEW EXERCISE 1

Select the correct answer in each of the following:

- (i) The order of matrix $\begin{bmatrix} 2 & 1 \end{bmatrix}$ is
 (a) 2-by-1 (b) 1-by-2
 (c) 1-by-1 (d) 2-by-2
- (ii) $\begin{bmatrix} \sqrt{2} & 0 \\ 0 & \sqrt{2} \end{bmatrix}$ is called _____ matrix
 (a) zero (b) unit
 (c) scalar (d) singular
- (iii) Which is order of square matrix
 (a) 2-by-2 (b) -by-2
 (c) 2-by-1 (d) 3-by-2
- (iv) Order of transpose of $\begin{bmatrix} 2 & 1 \\ 0 & 1 \\ 3 & 2 \end{bmatrix}$ is
 (a) 3-by-2 (b) 2-by-3
 (c) 1-by-3 (d) 3-by-1
- (v) Adjoint of $\begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$ is
 (a) $\begin{bmatrix} -1 & -2 \\ 0 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -2 \\ 0 & -1 \end{bmatrix}$
 (c) $\begin{bmatrix} -1 & 2 \\ 0 & -1 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}$
- (vi) Product of $\begin{bmatrix} x & y \\ 2 & -1 \end{bmatrix}$ is
 (a) $[2x + y]$ (b) $[x - 2y]$
 (c) $[2x - y]$ (d) $[x + 2y]$
- (vii) If $\begin{vmatrix} 2 & 6 \\ 3 & x \end{vmatrix} = 0$, then x is equal to
 (a) 9 (b) -6
 (c) 6 (d) -9
- (viii) If $X + \begin{bmatrix} -1 & -2 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ then x is equal to
 (a) $\begin{bmatrix} 2 & 2 \\ 2 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 0 & 2 \\ 2 & 2 \end{bmatrix}$
 (c) $\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ (d) $\begin{bmatrix} 2 & 2 \\ 0 & 2 \end{bmatrix}$

ANSWERS

(i)	b	(ii)	c	(iii)	a	(iv)	b
(v)	a	(vi)	c	(vii)	a	(viii)	d

Q.2: Complete the following:

- (i) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ is called _____ matrix.
- (ii) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ is called _____ matrix.
- (iii) Additive inverse of $\begin{bmatrix} 1 & -2 \\ 0 & -1 \end{bmatrix}$ is _____
- (iv) Matrix multiplication in general AB _____ BA.
- (v) Matrix A + B may be find if order of A and B is _____
- (vi) A matrix is called _____ matrix if number of rows and columns are equal.

ANSWERS

(i)	null	(ii)	Unit	(iii)	$\begin{bmatrix} -1 & 2 \\ 0 & 1 \end{bmatrix}$
(iv)	\neq	(v)	same	(vi)	square

Q.3: If $\begin{bmatrix} a+3 & 4 \\ 6 & b-1 \end{bmatrix} = \begin{bmatrix} -3 & 4 \\ 6 & 2 \end{bmatrix}$, then find a and b.

According to define of equal matrices.

Sol. $a + 3 = -3$
 $a = -3 - 3$
 $a = -6$
 $b - 1 = 2$
 $b = 2 + 1 \Rightarrow \mathbf{b = 3}$

Q.4: If $A = \begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 5 & -4 \\ -2 & -1 \end{bmatrix}$ then find the following:

- (i) $2A + 3B$ (ii) $-3A + 2B$
 (iii) $-3(A + 2B)$ (iv) $\frac{2}{3}(2A - 3B)$

(i) $2A + 3B$

Sol. Find $2A + 3B$
 $= 2\begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix} + 3\begin{bmatrix} 5 & -4 \\ -2 & -1 \end{bmatrix}$
 $= \begin{bmatrix} 4 & 6 \\ 2 & 0 \end{bmatrix} + \begin{bmatrix} 15 & -12 \\ -6 & -3 \end{bmatrix}$
 $= \begin{bmatrix} 4+15 & 6-12 \\ 2-6 & 0-3 \end{bmatrix}$
 $= \begin{bmatrix} 19 & -6 \\ -4 & -3 \end{bmatrix}$

(ii) $-3A + 2B$

$$\begin{aligned} \text{Sol. } &= -3 \begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix} + 2 \begin{bmatrix} 5 & -4 \\ -2 & -1 \end{bmatrix} \\ &= \begin{bmatrix} -6 & -9 \\ -3 & 0 \end{bmatrix} + \begin{bmatrix} 10 & -8 \\ -4 & -2 \end{bmatrix} \\ &= \begin{bmatrix} -6+10 & -9-8 \\ -3-4 & 0-2 \end{bmatrix} \\ &= \begin{bmatrix} 4 & -17 \\ -7 & -2 \end{bmatrix} \end{aligned}$$

(iii) $-3(A + 2B)$

$$\begin{aligned} \text{Sol. } -3(A + 2B) &= -3 \left(\begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix} + 2 \begin{bmatrix} 5 & -4 \\ -2 & -1 \end{bmatrix} \right) \\ &= -3 \left(\begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 10 & -8 \\ -4 & -2 \end{bmatrix} \right) \\ &= -3 \left(\begin{bmatrix} 2+10 & 3-8 \\ 1-4 & 0-2 \end{bmatrix} \right) \\ &= -3 \begin{bmatrix} 12 & -5 \\ -3 & -2 \end{bmatrix} \\ -3(A + 2B) &= \begin{bmatrix} -36 & 15 \\ 9 & 6 \end{bmatrix} \end{aligned}$$

(iv) $\frac{2}{3}(2A - 3B)$

$$\begin{aligned} \text{Sol. } &= \frac{2}{3} \left(2 \begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix} - 3 \begin{bmatrix} 5 & -4 \\ -2 & -1 \end{bmatrix} \right) \\ &= \frac{2}{3} \left(\begin{bmatrix} 4 & 6 \\ 2 & 0 \end{bmatrix} - \begin{bmatrix} 15 & -12 \\ -6 & -3 \end{bmatrix} \right) \\ &= \frac{2}{3} \left(\begin{bmatrix} 4-15 & 6+12 \\ 2+6 & 0+3 \end{bmatrix} \right) = \frac{2}{3} \begin{bmatrix} 11 & 18 \\ 8 & 3 \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{2}{3}\right)(-11) & \left(\frac{2}{3}\right)(18) \\ \left(\frac{2}{3}\right)(8) & \left(\frac{2}{3}\right)(3) \end{bmatrix} \\ &= \begin{bmatrix} \frac{-22}{3} & 12 \\ \frac{16}{3} & 2 \end{bmatrix} \end{aligned}$$

Q.5: Find the value of X, if $\begin{bmatrix} 2 & 1 \\ 3 & -3 \end{bmatrix} + X =$

$$\begin{aligned} &\begin{bmatrix} 4 & -2 \\ -1 & -2 \end{bmatrix} \\ \text{Sol. } &\begin{bmatrix} 2 & 1 \\ 3 & -3 \end{bmatrix} + X = \begin{bmatrix} 4 & -2 \\ -1 & -2 \end{bmatrix} \\ X &= \begin{bmatrix} 4 & -2 \\ -1 & -2 \end{bmatrix} - \begin{bmatrix} 2 & 1 \\ 3 & -3 \end{bmatrix} \end{aligned}$$

$$X = \begin{bmatrix} 4-2 & -2-1 \\ -1-3 & -2+3 \end{bmatrix}$$

$$X = \begin{bmatrix} 2 & -3 \\ -4 & 1 \end{bmatrix} \quad \text{Ans.}$$

Q.6: If $A = \begin{bmatrix} 0 & 1 \\ 2 & -3 \end{bmatrix}$, $B = \begin{bmatrix} -3 & 4 \\ 5 & -2 \end{bmatrix}$ then prove that:

(i) $AB \neq BA$ (ii) $A(BC) = (AB)C$

(i) $AB \neq BA$

$$\begin{aligned} \text{Sol. } AB &= \begin{bmatrix} 0 & 1 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} -3 & 4 \\ 5 & -2 \end{bmatrix} \\ &= \begin{bmatrix} (0)(-3)+(1)(5) & (0)(4)+(1)(-2) \\ (2)(-3)+(-3)(5) & (2)(4)+(-3)(-2) \end{bmatrix} \\ AB &= \begin{bmatrix} 0+5 & 0-2 \\ -6-15 & 8+6 \end{bmatrix} \dots\dots (i) \\ AB &= \begin{bmatrix} 5 & -2 \\ -21 & 14 \end{bmatrix} \\ BA &= \begin{bmatrix} -3 & 4 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & -3 \end{bmatrix} \\ &= \begin{bmatrix} (-3)(0)+(4)(2) & (-3)(1)+(4)(-3) \\ (5)(0)+(-2)(2) & (5)(1)+(-2)(-3) \end{bmatrix} \\ &= \begin{bmatrix} 0+8 & -3-12 \\ 0-4 & 5+6 \end{bmatrix} \\ BA &= \begin{bmatrix} 8 & -15 \\ -4 & 11 \end{bmatrix} \dots\dots (ii) \end{aligned}$$

From (i) and (ii) Its proved that $AB \neq BA$

Q.7: If $A = \begin{bmatrix} 3 & 2 \\ 1 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 4 \\ -3 & -5 \end{bmatrix}$ then

verify that: (i) $(AB)^t = B^t A^t$

(ii) $(AB)^{-1} = B^{-1} A^{-1}$

(i) $(AB)^t = B^t A^t$

Sol. $(AB)^t$

L.H.S.

$$\begin{aligned} AB &= \begin{bmatrix} 3 & 2 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 2 & 4 \\ -3 & -5 \end{bmatrix} \\ &= \begin{bmatrix} (3)(2)+(2)(-3) & (3)(4)+(2)(-5) \\ (1)(2)+(-1)(-3) & (1)(4)+(-1)(-5) \end{bmatrix} \\ AB &= \begin{bmatrix} 6-6 & 12-10 \\ 2+3 & 4+5 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 2 \\ 5 & 9 \end{bmatrix} \end{aligned}$$

Taking transport

$$(AB)^t = \begin{bmatrix} 0 & 2 \\ 5 & 9 \end{bmatrix} \\ = \begin{bmatrix} 0 & 5 \\ 2 & 9 \end{bmatrix}$$

R.H.S $B^t A^t$

$$B^t = \begin{bmatrix} 2 & -3 \\ 4 & -5 \end{bmatrix}, A^t = \begin{bmatrix} 3 & 1 \\ 2 & -1 \end{bmatrix},$$

$$B^t A^t = \begin{bmatrix} 2 & -3 \\ 4 & -5 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 2 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} (2)(3)+(-3)(2) & (2)(1)+(-3)(-1) \\ (4)(3)+(-5)(2) & (4)(1)+(-5)(-1) \end{bmatrix}$$

$$= \begin{bmatrix} 6-6 & 2+3 \\ 12-10 & 4+5 \end{bmatrix}$$

$$B^t A^t = \begin{bmatrix} 0 & 5 \\ 2 & 9 \end{bmatrix}$$

From (i) and (ii) its proved that $(AB)^t = B^t A^t$ **(ii)** $(AB)^{-1} = B^{-1} A^{-1}$ **Sol.** $(AB)^{-1}$

$$AB = \begin{bmatrix} 3 & 2 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 2 & 4 \\ -3 & -5 \end{bmatrix}$$

$$= \begin{bmatrix} (3)(2)+(2)(-3) & (3)(4)+(2)(-5) \\ (1)(2)+(-1)(-3) & (1)(4)+(-1)(-5) \end{bmatrix}$$

$$= \begin{bmatrix} 6-6 & 12-10 \\ 2+3 & 4+5 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 2 \\ 5 & 9 \end{bmatrix}$$

$$(AB)^{-1} = B^{-1} A^{-1}$$

We already have

$$AB = \begin{bmatrix} 0 & 2 \\ 5 & 9 \end{bmatrix}$$

$$|AB| = \begin{vmatrix} 0 & 2 \\ 5 & 9 \end{vmatrix} = (0)(9) - (2)(5)$$

$$|AB| = 0 - 10 = -10 \neq 0 \text{ Non-singular matrix}$$

so solution possible.

$$\text{Adj } AB = \begin{bmatrix} 9 & -2 \\ -5 & 0 \end{bmatrix}$$

$$(AB)^{-1} = \frac{1}{|AB|} \text{Adj } AB$$

$$= -\frac{1}{10} \begin{bmatrix} 9 & -2 \\ -5 & 0 \end{bmatrix} x$$

$$\text{As } B^{-1} = \frac{1}{|B|} \text{Adj } B$$

$$|B| = \begin{vmatrix} 2 & 4 \\ -3 & -5 \end{vmatrix} = (2)(-5) - (-3)(4)$$

$$= -10 + 12 = 2$$

$$B^{-1} = \frac{1}{2} \begin{bmatrix} -5 & -4 \\ +3 & 2 \end{bmatrix} \quad \dots(i)$$

$$\text{As } A^{-1} = \frac{1}{|A|} \text{Adj } A$$

$$|A| = \begin{vmatrix} 3 & 2 \\ 1 & -1 \end{vmatrix} = (3)(-1) - (1)(2) = -3 - 2$$

$$= -5$$

$$A^{-1} = \frac{1}{-5} \begin{bmatrix} -1 & -2 \\ -1 & 3 \end{bmatrix} \quad x$$

Multiply (i) \times (ii)

$$B^{-1} \times A^{-1} = \frac{1}{2} \begin{bmatrix} -5 & -4 \\ 3 & 2 \end{bmatrix} \times \frac{1}{-5} \begin{bmatrix} -1 & -2 \\ -1 & 3 \end{bmatrix}$$

$$= \frac{1}{2} \times \frac{1}{-5} \begin{bmatrix} -5 & -4 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} -1 & -2 \\ -1 & 3 \end{bmatrix}$$

$$= \frac{-1}{10} \begin{bmatrix} (-5)(-1)+(-1)(-4) & (-5)(-2)+(-4)(3) \\ (3)(-1)+(-1)(-4) & (3)(-2)+(-1)(3) \end{bmatrix}$$

$$= \frac{-1}{10} \begin{bmatrix} 5+4 & 10-12 \\ -3-2 & -6+6 \end{bmatrix}$$

$$= \frac{-1}{10} \begin{bmatrix} 9 & -2 \\ -5 & 0 \end{bmatrix} = y$$

See equation x and y $(AB)^{-1} = B^{-1} \times A^{-1}$

IMPORTANT KEY POINTS

OF CHAPTER

- A rectangular array of real numbers enclosed with brackets is said to form a matrix.
- A matrix A is called rectangular, if the number of rows and number of columns of A are not equal.
- A matrix A is called a square matrix, if the number of rows of A is equal to the number of columns.
- A matrix A is called a row matrix, if A has only one row.
- A matrix A is called a column matrix, if A has only one column.
- A matrix A is called a null or zero matrix, if each of its entry is 0.
- Let A be a matrix. The matrix A^t is a new matrix which is called transpose of matrix A and is obtained by interchanging rows of A into its respective columns (or columns into respective rows).
- A square matrix A is called symmetric, if $A^t = A$.

- Let A be a matrix. Then its negative, -A, is obtained by changing
- The signs of all the non - zero entries of A.
- A square matrix M is said to be skew symmetric, if $M^t = -M$,
- A square matrix M is called a diagonal matrix, if atleast any one of entry of its diagonal is not zero and all non diagonal entries are zero.
- A scalar matrix is called identity matrix, if all diagonal entries are 1.

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ is called 3-by-3 identity matrix,}$$

- Let A be a matrix of order 2-by-3. Then a matrix B of same order is said to be an additive identity of matrix A, if $B + A = A = A + B$
- Let A be a matrix. A matrix B is defined as an additive inverse of A, if $B + A = O = A + B$
- Let A be a matrix. Another matrix B is called the identity matrix of A under multiplication, if $B \times A = A = A \times B$.

Let $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be a 2-by-2 matrix. A real number is called determinant of M, denoted by $\det M$ such that

$$\det M = \begin{vmatrix} a & c \\ b & d \end{vmatrix} = ad - bc = \lambda$$

- A square matrix M is called singular, if the determinant of M is equal to zero.
- A square matrix M is called non-singular, if the determinant of M is not equal to zero.
- For a matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, adjoint of M is defined by

$$\text{Adj } M = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

➤ Let M be a square matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then

$$M^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \text{ where,}$$

$$\det M = \begin{vmatrix} a & c \\ b & d \end{vmatrix} = ad - bc = \lambda \neq 0$$

- Any two matrices A and B are called equal, if order of A = order of B (ii) corresponding entries are equal
- Any two matrices M and N are said to be conformable for addition, if order of M = order of N.
- The following laws of addition hold $M + N = N + M$ (Commutative)
 $(M + N) + T = M + (N + T)$ (Associative)
- The matrices M and N are conformable for multiplication to obtain MN if the number of columns of M = number of rows of N, where
- (i) $(MN) \neq NM$, in general
- (ii) $(MN)T = M(NT)$ (Associative law)
- (iii) $M(N + T) = MN + MT$
- (iv) $(N + T)M = NM + TM$ } (Distributive laws)
- Law of transpose of product $(AB)^t = B^t A^t$
 $AA^{-1} = I = A^{-1}A$
- The solution of a linear system of equations,

$$ax + by = m$$

$$cx + dy = n$$

by expressing in matrix form

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} m \\ n \end{bmatrix} \text{ is given}$$

$$\text{by } \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} \begin{bmatrix} m \\ n \end{bmatrix} \text{ if the}$$

coefficient matrix is non-singular.

- By using the Cramer's rule the determinantal form of the solution is

$$x = \frac{\begin{vmatrix} m & b \\ n & d \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}} \text{ and } y = \frac{\begin{vmatrix} a & m \\ c & n \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}}$$

$$\text{Where } \begin{vmatrix} a & b \\ c & d \end{vmatrix} \neq 0$$

ADDITIONAL MCQ'S

1. Adjoint of $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is
- (a) $\begin{bmatrix} a & -b \\ -c & d \end{bmatrix}$ (b) $\begin{bmatrix} -a & b \\ c & -d \end{bmatrix}$
- (c) $\begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ (d) $\begin{bmatrix} a & c \\ b & d \end{bmatrix}$
2. The idea of matrices was given by
- (a) Aurthur Cayley (b) Briggs
(c) Al-Khwarzmi (d) Thomas Harriot
3. $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then $|A| = \dots\dots$
- (a) $ab-cd$ (b) $ac-bd$
(c) $bc-ad$ (d) $ad-bc$
4. Aurthur Cayley introduced theory of matrices in
- (a) 1854 (b) 1856
(c) 1858 (d) 1860
5. For
- values of $x \begin{bmatrix} 3 & -6 \\ 2 & x \end{bmatrix}$ will be a singular matrix.
- (a) -3 (b) -4
(c) 3 (d) 4
6. Product of $\begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ -4 \end{bmatrix}$.
- (a) [-13] (b) [-3]
(c) [3] (d) [13]
7. If $\begin{bmatrix} a+3 & 4 \\ 6 & 0 \end{bmatrix} = \begin{bmatrix} -3 & 4 \\ 6 & 0 \end{bmatrix}$ value of 'a' will be.
- (a) 6 (b) 3
(c) -3 (d) -6
8. The order of matrix $\begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$.
- (a) 1-by-3 (b) 3-by-1
(c) 3-by-3 (d) 2-by-2
9. A square matrix M is called to be skew symmetric,
- (a) $M^t = \overline{M}$ (b) $M^t = \frac{1}{M}$
(c) $M^t = -M$ (d) $M^t = M$

10. A square matrix M is called to be symmetric matrix if:

- (a) $M^t = \overline{M}$ (b) $M^t = \frac{1}{M}$
(c) $M^t = -M$ (d) $M^t = M$

ANSWERS

(i)	c	(ii)	a	(iii)	d	(iv)	c
(v)	b	(vi)	b	(vii)	d	(viii)	b
(ix)	c	(x)	d				